

Mark Scheme (Results)

June 2011

International GCSE

Mathematics (4PM0) Paper 02

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.


Ask The Expert can be accessed online at the following link:
<http://www.edexcel.com/Aboutus/contact-us/>

June 2011

Publications Code UG027963

All the material in this publication is copyright

© Edexcel Ltd 2011

Q.	Scheme	Marks	
1.	$\sum = \frac{15}{2}(9+37), = 345$	M1A1,A1 (3)	
2.	$v = 3t^2 + 4t - 3$ $12 = 3t^2 + 4t - 3$ $0 = 3t^2 + 4t - 15$ $0 = (3t-5)(t+3)$ $t = \frac{5}{3}$	M1A1 M1 A1 (4)	
3.		(a) $\frac{\sin C}{5} = \frac{\sin 25}{3}$ $\sin C = \frac{5 \sin 25}{3}$ $C = 135.2\dots = 135$ (b) $CD = 2 \times 3 \cos 44.8$ $CD = 4.257\dots = 4.26 \text{ cm}$	M1A1 A1 M1A1ft A1 (6)
4.	(a) $x = -4 \quad 0 = (-4)^3 + 2(-4)^2 - 11(-4) - m$ $m = -64 + 32 + 44 = 12 \quad *$ (b) $x^3 + 2x^2 - 11x - 12 = (x+4)(x^2 - 2x - 3)$ $= (x+4)(x-3)(x+1)$ (c) $b = -1 \quad d = 3$	M1 A1 cs B1 M1A1 B1 (6)	
5.	(a) $\frac{5 \times 1 + 2q}{3} = 13 \quad 2q = 39 - 5 \quad q = 17$ $\frac{p + 2 \times 12}{3} = 10 \quad p = 30 - 24 = 6$ (b) $A \text{ --- } 3 \text{ --- } B \text{ --- } 2 \text{ --- } E$ $17\mathbf{i} + 12\mathbf{j} = \frac{2(5\mathbf{i} + 6\mathbf{j}) + 3\mathbf{e}}{5}$ $85\mathbf{i} + 60\mathbf{j} = 10\mathbf{i} + 12\mathbf{j} + 3\mathbf{e}$ $3\mathbf{e} = 75\mathbf{i} + 48\mathbf{j}$ $\mathbf{e} = 25\mathbf{i} + 16\mathbf{j}$	M1A1 A1 M1 A1 A1 (6)	

Q.	Scheme	Marks
6	<p>(a) $8\theta = 6 \quad \theta = \frac{3}{4}$ (accept 0.75) oe</p> <p>(b) $\frac{1}{2}r^2\theta = \frac{1}{2} \times 8^2 \times \frac{3}{4} = 24 \text{ cm}^2$</p> <p>(c) Area of $\Delta ABC = \frac{1}{2} \times 8^2 \times \sin AOB = 21.81\dots$</p> <p>Area of segment $= 24 - 21.81\dots = 2.187 = 2.19 \text{ cm}^2$</p>	M1A1 M1A1 M1A1 A1ft (7)
7.	<p>(a) $V = 3x^2h = 30$ $S = 3x^2 + 2xh + 2 \times 3xh$ $xh = \frac{10}{x} \Rightarrow S = 3x^2 + 2 \times \frac{10}{x} + 6 \times \frac{10}{x}$ $S = 3x^2 + \frac{80}{x} *$</p> <p>(b) $\frac{dS}{dx} = 6x - \frac{80}{x^2}$ $\frac{dS}{dx} = 0 \quad 6x^3 = 80 \quad x = \sqrt[3]{\frac{40}{3}} \quad (= 2.371\dots)$ $S_{\min} = 3\left(\sqrt[3]{\frac{40}{3}}\right)^2 + \frac{80}{\sqrt[3]{40}} = 50.60\dots = 50.6 \text{ cm}^3$</p> <p>(c) $\frac{d^2S}{dx^2} = 6 + \frac{160}{x^3} > 0 \text{ for } x > 0$ $\therefore \text{minimum}$</p>	B1 B1 M1 A1 M1 M1A1 M1A1 M1 A1ft (11)

Q.	Scheme	Marks
8.	<p>(a) $a + ar^2 = 100, \quad ar + ar^2 = 60$</p> $\frac{1+r^2}{r+r^2} = \frac{100}{60}$ $6+6r^2 = 10r+10r^2 \quad 2r^2 + 5r - 3 = 0$ $(2r-1)(r+3) = 0$ $r = \frac{1}{2} \quad r = -3$ <p>(b) $r = \frac{1}{2} \quad a = \frac{100}{1+(\frac{1}{2})^2} = 80$</p> <p>(c) $S_n = \frac{a(1-r^n)}{1-r} = \frac{80\left(1-\left(\frac{1}{2}\right)^n\right)}{1-\frac{1}{2}} > 159.9$</p> $\frac{159.9}{160} < 1 - \left(\frac{1}{2}\right)^n$ $\left(\frac{1}{2}\right)^n = 1 - \frac{159.9}{160}$ $n \log 0.5 < \log\left(1 - \frac{159.9}{160}\right)$ $n > \frac{\log\left(1 - \frac{159.9}{160}\right)}{\log 0.5} = 10.6$ <p>n=11</p>	<p>M1,A1</p> <p>M1</p> <p>A1A1</p> <p>M1A1</p> <p>M1A1</p> <p>M1</p> <p>A1 (11)</p>

Q.	Scheme	Marks
9	<p>(a) $\left(1 - \frac{3x}{4}\right)^{\frac{1}{3}} = 1 + \left(\frac{1}{3}\right)\left(-\frac{3x}{4}\right) + \frac{\left(\frac{1}{3}\right)\left(-\frac{2}{3}\right)}{2!} \left(-\frac{3x}{4}\right)^2 + \frac{\left(\frac{1}{3}\right)\left(-\frac{2}{3}\right)\left(-\frac{5}{3}\right)}{3!} \left(-\frac{3x}{4}\right)^3$ $= 1 - \frac{x}{4} - \frac{x^2}{16} - \frac{5x^3}{192}$</p> <p>(b) $\left(1 + \frac{3x}{4}\right)^{-\frac{1}{3}} = 1 + \left(-\frac{1}{3}\right)\left(\frac{3x}{4}\right) + \frac{\left(-\frac{1}{3}\right)\left(-\frac{4}{3}\right)}{2!} \left(\frac{3x}{4}\right)^2 + \frac{\left(-\frac{1}{3}\right)\left(-\frac{4}{3}\right)\left(-\frac{7}{3}\right)}{3!} \left(\frac{3x}{4}\right)^3$ $= 1 - \frac{x}{4} + \frac{x^2}{8} - \frac{7x^3}{96}$</p> <p>(c) $x < \frac{4}{3}$</p> <p>(d) $\left(\frac{4-3x}{4+3x}\right)^{\frac{1}{3}} = \left(\frac{1-\frac{3x}{4}}{1+\frac{3x}{4}}\right)^{\frac{1}{3}} = \left(1 - \frac{x}{4} - \frac{x^2}{16} - \frac{5x^3}{192}\right) \left(1 - \frac{x}{4} + \frac{x^2}{8} - \frac{7x^3}{96}\right)$ $= 1 - \frac{x}{4} + \frac{x^2}{8} - \frac{7x^3}{96} - \frac{x}{4} + \frac{x^2}{16} - \frac{x^3}{32} - \frac{x^2}{16} + \frac{x^3}{64} - \frac{5x^3}{192}$ $= 1 - \frac{x}{2} + \frac{x^2}{8} - \frac{11x^3}{96}$</p> <p>(e) $\int_0^{0.5} \left(\frac{4-3x}{4+3x}\right)^{\frac{1}{3}} dx = \int_0^{0.5} \left(1 - \frac{x}{2} + \frac{x^2}{8} - \frac{11x^3}{96}\right) dx$ $= \left[x - \frac{x^2}{4} + \frac{x^3}{24} - \frac{11x^4}{384} \right]_0^{0.5}$ $= 0.5 - \frac{0.5^2}{4} + \frac{0.5^3}{24} - \frac{11 \times 0.5^4}{384} = 0.4409\dots = 0.441$</p>	M1 A1A1 M1 A1A! B1 M1 M1 A1 M1A1ft M1A1 (14)
10	<p>(a) $\alpha + \beta = -6 \quad \alpha\beta = 2$</p> <p>(i) $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta = 36 - 4 = 32$</p> <p>(ii) $\alpha^4 + \beta^4 = (\alpha^2 + \beta^2)^2 - 2(\alpha\beta)^2 = 32^2 - 8 = 1016$</p> <p>(b) $(\alpha - \beta)^2 = \alpha^2 - 2\alpha\beta + \beta^2 = 32 - 4 = 28$ $\alpha - \beta = \sqrt{28} = 2\sqrt{7}$</p> <p>(c) $\alpha^4 - \beta^4 = (\alpha^2 + \beta^2)(\alpha^2 - \beta^2), = (\alpha^2 + \beta^2)(\alpha + \beta)(\alpha - \beta)$</p> <p>(d) $\alpha^4 - \beta^4 = 32 \times (-6) \times 2\sqrt{7} = -384\sqrt{7}$</p> <p>(e) $(\alpha^4 + \beta^4) - (\alpha^4 - \beta^4) = 2\beta^4$ $2\beta^4 = 1016 + 384\sqrt{7}$ $\beta^4 = 508 + 192\sqrt{7}$</p>	B1 M1A1 M1A1 M1A1 A1 M1A1 M1A1 M1 A1 M1 A1 (14)

Q.	Scheme	Marks
11.	<p>(a) $x^2 + 6x + 8 = (x+3)^2 - 9 + 8 = (x+3)^2 - 1$</p> <p>(b) $f(x)$ is least when $x = -3$ least value is -1</p> <p>(c) $x^2 + 6x + 8 = 2 - x$ $x^2 + 7x + 6 = 0$ $(x+6)(x+1) = 0$ $x = -6 \quad x = -1$</p> <p>(d) $x^2 + 6x + 8 = 0$ $(x+2)(x+4) = 0$ $x = -2 \quad x = -4$</p> <p>(e) </p> <p>(f) Area = $\int_{-6}^{-1} \{2 - x - (x^2 + 6x + 8)\} dx$ $= \left[-\frac{x^3}{3} - \frac{7x^2}{2} - 6x \right]_{-6}^{-1}$ $= \left(\frac{1}{3} - \frac{7}{2} + 6 \right) - \left(\frac{6^3}{3} - \frac{7 \times 6^2}{2} + 6^2 \right)$ $= 20\frac{5}{6}$ or awrt 20.8</p>	M1A1A1 B1 B1 M1 M1 A1A1 M1A1 B1 B1 M1 M1A1 M1 M1A1 M1 M1A1 M1 M1 A1 (18)

Further copies of this publication are available from
International Regional Offices at www.edexcel.com/international



For more information on Edexcel qualifications, please visit
www.edexcel.com

Alternatively, you can contact Customer Services at
www.edexcel.com/ask or on + 44 1204 770 696



Llywodraeth Cynulliad Cymru
Welsh Assembly Government

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

