

Mark Scheme (Results)

Summer 2013

GCE Mechanics 4 (6680/01)

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2013

Publications Code UA036433

All the material in this publication is copyright

© Pearson Education Ltd 2013

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL GCE MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes:

- bod – benefit of doubt
 - ft – follow through
 - the symbol \surd will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
 7. Ignore wrong working or incorrect statements following a correct answer.
 8. In some instances, the mark distributions (e.g. M1, B1 and A1) printed on the candidate's response may differ from the final mark scheme


General Rules for Marking Mechanics



- Usual rules for M marks: correct no. of terms; dim correct; all terms that need resolving (i.e. multiplied by cos or sin) are resolved.
- Omission or extra g in a resolution is accuracy error not method error.
- Omission of mass from a resolution is method error.
- Omission of a length from a moments equation is a method error.
- Omission of units or incorrect units is not (usually) counted as an accuracy error.
- DM indicates a dependent method mark i.e. one that can only be awarded if a previous specified method mark has been awarded.
- Any numerical answer which comes from use of $g = 9.8$ should be given to 2 or 3 SF.
- Use of $g = 9.81$ should be penalised once per (complete) question.
- N.B. Over-accuracy or under-accuracy of correct answers should only be penalised *ONCE* per complete question.
- In all cases, if the candidate clearly labels their working under a particular part of a question i.e. (a) or (b) or (c),.....then that working can only score marks for that part of the question.
- Accept column vectors in all cases.

Misreads – if a misread does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, bearing in mind that after a misread, the subsequent A marks affected are treated as A ft.

Question Number	Scheme	Marks	
1(a)	Equation of motion: $\frac{1}{2}g - \frac{3}{2}v = \frac{1}{2} \frac{dv}{dt}$	M1 A1	Differential equation. All 3 terms required but condone sign errors
NB: these two marks are available in (b) if not scored in (a)			
	$\int 1 dt = \int \frac{1}{9.8 - 3v} dv$ $t + (C) = -\frac{1}{3} \ln(9.8 - 3v)$ $t = 0, v = 0 \Rightarrow C = -\frac{1}{3} \ln 9.8$ $t = -\frac{1}{3} \ln\left(\frac{9.8 - 3v}{9.8}\right)$ $3v = 9.8(1 - e^{-3t})$ *Given Answer*	M1 A1=A1 M1 A1 A1 (8)	Separate the variables and attempt to integrate A1 for each side. C not needed Use initial conditions to evaluate C or limits on a definite integral. Or equivalent Watch out. cwo
(a) alt	Equation of motion: $\frac{1}{2}g - \frac{3}{2}v = \frac{1}{2} \frac{dv}{dt}$ $e^{3t} \frac{dv}{dt} + 3e^{3t}v = ge^{3t}, \frac{d}{dt}(ve^{3t}) = ge^{3t}$ $ve^{3t} = \frac{1}{3}ge^{3t} (+c)$ $t = 0, v = 0 \Rightarrow 0 = \frac{1}{3}g + C$ $\Rightarrow ve^{3t} = \frac{1}{3}g(e^{3t} - 1), 3v = 9.8(1 - e^{-3t})$	M1 A1 M1 A1=A1 M1 A1 A1	All 3 terms required but condone sign errors Use of integrating factor e^{3t} A1 for each side. +C not required. Use initial conditions to evaluate C Correct equation in any equivalent form Given form cwo


Question Number	Scheme	Marks	
1(b)	$\frac{dx}{dt} = \frac{9.8}{3}(1 - e^{-3t}) \Rightarrow x = \frac{9.8}{3}\left(t + \frac{1}{3}e^{-3t}\right) + C$ $t = 0, x = 0 \Rightarrow C = -\frac{9.8}{9}$ $t = 2, x \approx 5.4 \text{ (m)}$	M1 A1 M1 A1 A1 (5) (13)	Integrate the given v to find x C not needed Use the initial conditions to evaluate C or use limits correctly in a definite integral 5.45, $\frac{g}{9}(5 + e^{-6})$ or equivalent
(b) alt	$g - 3v = v \frac{dv}{dx}$		
	$\int 1 dx = \int \frac{v}{g - 3v} dv = \int -\frac{1}{3} + \frac{g}{3(g - 3v)} dv$	M1	Separate the variables and rearrange the RHS
	$x = -\frac{v}{3} - \frac{g}{9} \ln(g - 3v) + C$	A1	+ C not needed
	$x = 0, v = 0 \Rightarrow C = \frac{g}{9} \ln g \text{ and}$ $t = 2, v = \frac{g}{3}(1 - e^{-6}) (= 3.258\dots)$	M1 A1	Use the initial conditions to find C & find the value of v when $t = 2$
	$x = \frac{g}{9}(1 - e^{-6}) - \frac{g}{9} \ln(e^{-6}) = 5.4$	A1	
		(5)	
		(13)	



Question Number	Scheme	Marks	
2(a)	Shortest time $50 \div \frac{10}{9} = 45$ (s)	M1,A1	
(b)	Drifts $\frac{2}{3} \times 45 = 30$ (m)	M1 A1	$\frac{2}{3} \times \text{their time}$
(c)	 <p data-bbox="705 343 1008 454">Trig or pythag to find velocity of swimmer in direction AB</p> <p data-bbox="705 550 795 622">$\frac{8}{9} \text{ms}^{-1}$</p> <p data-bbox="414 790 660 853">$50 \div \frac{8}{9} = 56.25$ (s)</p>	M1 A1 DM1,A1 (8)	0.88 or better Dependent on the previous M 56 or better


Question Number	Scheme	Marks	
3	 <p> $0.6u$ or $u \cos \alpha$ $1.2u$ or $2u \cos \alpha$ $2m \times 1.2u - 3m \times 0.6u = 3ma + 2mb$ $(3a + 2b = 0.6u)$ $e(1.2u + 0.6u) = a - b$ $(a - b = 0.3u)$ $a = 0.24u$ or $b = -0.06u$ $(1.2u - (-0.06u)) \times 2m = 2.52mu$ or $(0.24u - (-0.6u)) \times 3m = 2.52mu$ </p>	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1ft</p> <p>M1</p> <p>A1ft</p> <p>DM1</p> <p>A1</p> <p>M1</p> <p>A1 (10)</p>	 <p> component of the initial velocity of A parallel to the line of centres on impact component of the initial velocity of B parallel to the line of centres on impact CLM parallel to the line of centres. Requires all the terms. Correct unsimplified for their $0.6u$ and $1.2u$ Restitution parallel to the line of centres. Must be used the right way round. Correct unsimplified for their $0.6u$ and $1.2u$ If signs are inconsistent between the two equations, penalise here. Solve a pair of simultaneous eqns in a & b for one of a & b. Dependent on the two previous M marks. In terms of u only Find impulse on A or B. Unsimplified. For their a or b. Correct mass for the velocities used. </p> <p>$\frac{63}{25}$</p>

Question Number	Scheme	Marks	
4(a)	PE of ring = $-mgx$ PE of particle = $-3mg(L - \sqrt{x^2 + d^2})$ $\Rightarrow V = 3mg\sqrt{x^2 + d^2} - mgx + \text{constant. AG}$	B1 M1 A1 A1	Taking the level of the peg as zero PE Watch out
(b)	$\frac{dV}{dx} = \frac{3mg \cdot 2x}{2\sqrt{x^2 + d^2}} - mg$ $\frac{dV}{dx} = 0 \Rightarrow 3x = \sqrt{x^2 + d^2}, 9x^2 = x^2 + d^2, 8x^2 = d^2$ $x = \frac{d}{\sqrt{8}} = \left(\frac{\sqrt{2}d}{4}\right)$	M1 M1 A1	(4) Set $\frac{dV}{dx} = 0$ and solve for x 0.354d of better
(c)	$\frac{d^2V}{dx^2} = 3mg \left(\frac{\sqrt{x^2 + d^2} \cdot 1 - x \cdot \frac{2x}{2\sqrt{x^2 + d^2}}}{x^2 + d^2} \right) =$ $3mg \left(\frac{\sqrt{9x^2} \cdot 1 - x \cdot \frac{2x}{2\sqrt{9x^2}}}{9x^2} \right) = \frac{3mgd^2}{(x^2 + d^2)^{\frac{3}{2}}} (> 0)$ Stable	M1 A1 A1ft	Product or quotient rule $\frac{d^2V}{dx^2} = \frac{3mg}{\sqrt{x^2 + d^2}} - \frac{3mgx}{2} \cdot 2x \cdot (x^2 + d^2)^{-\frac{3}{2}}$ OR $= 3mg \left(\frac{3x - \frac{x}{3}}{9x^2} \right) (> 0)$ Correct unsimplified. $\frac{16\sqrt{2}mg}{9d}, 2.5 \frac{mg}{d}, \frac{d^2V}{d\theta^2} = \frac{9mgd}{\sqrt{8}}$ Correct conclusion for their expression


(10)

Question Number	Scheme	Marks	
5(a)	 <p data-bbox="779 220 969 292">Minimum $V = 12 \cos 50^\circ$</p> <p data-bbox="779 411 869 443">≈ 7.71</p>	<p data-bbox="1126 220 1182 252">M1</p> <p data-bbox="1126 331 1182 363">A1</p> <p data-bbox="1126 411 1182 443">A1</p>	<p data-bbox="1261 220 1953 308">Use of triangle with right angle between v_c and $c v_s$. Condone sin/cos confusion. Correct unsimplified trig expression</p> <p data-bbox="1261 411 1395 451">7.71 only</p>

Question Number	Scheme	Marks	
5(b)	 <p data-bbox="465 1002 1079 1145">Vector triangle for relative velocities when $V = 14$ Select the vector triangle with the relative velocity due N.</p> $\frac{\sin \theta}{12} = \frac{\sin 40}{14}$ <p data-bbox="465 1273 645 1305">Bearing 033°</p>	<p data-bbox="1131 1002 1176 1034">M1</p> <p data-bbox="1131 1082 1176 1114">A1</p> <p data-bbox="1131 1193 1198 1225">DM1</p> <p data-bbox="1131 1233 1176 1265">A1</p> <p data-bbox="1131 1273 1176 1305">A1</p> <p data-bbox="1198 1313 1243 1345">(8)</p>	 <p data-bbox="1261 1002 1944 1074">could have relative velocity due S. Could show both possibilities.</p> <p data-bbox="1261 1193 1765 1225">Use of sine rule or equivalent to find θ</p> <p data-bbox="1261 1273 1630 1305">Final answer. Accept 33.4°</p>

Question Number	Scheme	Marks	
<p>6(a)</p>	 <p> $a + Ut = y + (a + x)$ $Ut = x + y$ *Answer Given* </p>	<p>M1 A1</p>	<p>Diagram or clear explanation using distances Watch out for fudges.</p>
<p>(b)</p>	<p> $T = \frac{9ma \times x}{a} = 9mx$ $T - 6m\ddot{y} = m\ddot{y}$ $9mx - 6m(U - \dot{x}) = -m\ddot{x}$ $\ddot{x} + 6\dot{x} + 9x = 6U$ </p>	<p>B1 M1 A2 A1</p>	<p>Equation of motion of P. Requires all 3 terms in terms of x and/or y Expressed in terms of x. -1 each error Answer given. Watch out for fudges</p>
<p>(c)</p>	<p> $t = 0, x = 0, \dot{x} = U \quad 0 = AU + \frac{2U}{3}, A = -\frac{2}{3}$ $\dot{x} = BUe^{-3t} - 3(A + Bt)Ue^{-3t}$ $U = BU - 3AU, B = 3A + 1 = -1$ </p>	<p>M1 A1 M1 A1 A1</p>	<p>Use initial conditions to find A Differentiate</p>
<p>(d)</p>	<p> $\dot{y} = U - \dot{x} = U - (-Ue^{-3t} + 2Ue^{-3t} + 3Ute^{-3t})$ $= U(1 - e^{-3t} - 3te^{-3t})$ </p>	<p>M1 A1</p>	<p>Or equivalent</p>
		(14)	

Question Number	Scheme	Marks	
7(a)	<p>State that impulse acts perpendicular to the wall and demonstrate that $(2\mathbf{i} + \mathbf{j}) \cdot (-\mathbf{i} + 2\mathbf{j}) = 0$</p>	B1	Requires scalar product or gradient diagram.
(b)	<p>Impulse momentum equation: $m(\mathbf{v} - \mathbf{u}) = m[(a-b)\mathbf{i} + a\mathbf{j}] = \lambda(-\mathbf{i} + 2\mathbf{j})$ $\Rightarrow a = -2(a - b), 3a = 2b$</p> <p>OR</p> <p>Taking scalar products of velocities with $(2\mathbf{i} + \mathbf{j})$ $\begin{pmatrix} b \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 1 \end{pmatrix} = 2b$ and $\begin{pmatrix} a \\ a \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 1 \end{pmatrix} = 3a$ No change parallel to the wall so $2b = 3a$.</p> <p>Scalar products with $(-\mathbf{i} + 2\mathbf{j})$: $\begin{pmatrix} b \\ 0 \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 2 \end{pmatrix} = -b$ and $\begin{pmatrix} a \\ a \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 2 \end{pmatrix} = a$</p> <p>Impact equation: $a = eb$ $e = \frac{2}{3}$</p>	M1 A2 A1 M1 A1A1 A1 B1 M1A1 A1	Requires all terms present and of the correct structure -1 each error

Question Number	Scheme	Marks	
7(b) alt	 <p> $b \cos \theta = a\sqrt{2} \cos(45 - \theta)$ $b \cos \theta = a \cos \theta + a \sin \theta, \quad 2b - 2a = a$ $2b = 3a$ Use of $\tan \theta = \frac{1}{2}$ $a\sqrt{2} \sin(45 - \theta) = eb \sin \theta$ $a \cos \theta = (a + eb) \sin \theta, \quad 2a = a + eb$ $e = \frac{2}{3}$ </p>	<p>M1 A2 A1 B1 M1 A1 A1</p>	<p>Parallel to the wall. Condone trig confusion? -1 each error. Both angles in same variable?</p> <p>When seen in (b). Implied by 26.6 or 18.4</p> <p>Perpendicular to the wall. Condone consistent trig confusion? $e = \sqrt{\frac{10a^2}{b^2} - 4}$</p> <p>0.67 or better</p>
(c)	<p>Fraction of KE lost = $\frac{b^2 - 2a^2}{b^2}$</p> $= \frac{1 - 2 \times \frac{4}{9}}{1} = \frac{1}{9}$	<p>M1A1 A1</p> <p>(12)</p>	

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email publication.orders@edexcel.com

Order Code UA036433 Summer 2013

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual



Llywodraeth Cynulliad Cymru
Welsh Assembly Government

