

Copyright © mppe.org.uk and its license. All Rights Reserved

**NOTICE TO CUSTOMER:**

**The sale of this product is intended for use of the original purchaser only and for use only on a single computer system. Duplicating, selling, or otherwise distributing this product is a violation of the law ; **your license of the product will be terminated at any moment if you are selling or distributing the products.****

No parts of this book may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher.

## mock papers 1

## SECTION A

1 In the reaction of manganate(VII) ions with reducing agents in strongly acidic solution, the half-reaction for the reduction is

- ☐ A  $\text{MnO}_4^- + 4\text{H}^+ + 3\text{e}^- \rightarrow \text{MnO}_2 + 2\text{H}_2\text{O}$
- ☐ B  $\text{MnO}_4^- + 4\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 2\text{H}_2\text{O}$
- ☐ C  $\text{MnO}_4^- + 8\text{H}^+ + 3\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$
- ☐ D  $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$

(Total for Question 1 = 1 mark)

2 In the titration of iodine with standard sodium thiosulfate solution, starch is often used as an indicator. The starch should **not** be added until nearly all the iodine has reacted because

- ☐ A it is decomposed by high concentrations of iodine.
- ☐ B the blue complex formed is bleached by high concentrations of iodine.
- ☐ C the blue complex formed with high concentrations of iodine is insoluble and does not re-dissolve as more thiosulfate is added.
- ☐ D the starch reacts with the thiosulfate ions being added.

(Total for Question 2 = 1 mark)

3 The conditions needed for the  $E^\ominus$  value of the standard hydrogen electrode to be exactly 0 V are

- ☐ A 1 mol dm<sup>-3</sup> solution of hydrogen ions, 1 atm pressure of hydrogen, 25°C.
- ☐ B 1 mol dm<sup>-3</sup> solution of hydrogen ions, 1 atm pressure of hydrogen, room temperature.
- ☐ C 1 mol dm<sup>-3</sup> solution of hydrogen ions, laboratory pressure of hydrogen, 25°C.
- ☐ D 0.1 mol dm<sup>-3</sup> solution of hydrogen ions, 1 atm pressure of hydrogen, 25°C.

(Total for Question 3 = 1 mark)

4 The electrode potential for a cell can be used to calculate the equilibrium constant for the cell reaction. This is because

- ☐ A  $E_{\text{cell}}^{\ominus}$  is proportional to  $\ln K$ .
- ☐ B  $E_{\text{cell}}^{\ominus}$  is proportional to  $K$ .
- ☐ C  $\ln E_{\text{cell}}^{\ominus}$  is proportional to  $\ln K$ .
- ☐ D  $\ln E_{\text{cell}}^{\ominus}$  is proportional to  $K$ .

(Total for Question 4 = 1 mark)

5 Which of the following successive ionization energies (values in  $\text{kJ mol}^{-1}$ ) could have come from a transition element?

- ☐ A 496 4563 6913 9544 13352 16611 20115 25941
- ☐ B 590 1145 4912 6474 8144 10496 12320 14207
- ☐ C 717 1509 3249 4940 6985 9200 11508 18956
- ☐ D 2081 3952 6122 9370 12177 15239 19999 23069

(Total for Question 5 = 1 mark)

6 Although platinum is very unreactive, it is used as a catalyst in catalytic converters in motor cars. Which of the following is true?

- ☐ A It converts nitrogen oxides and carbon monoxide to nitrogen and carbon dioxide by adsorbing the reactants on its surface so weakening their bonds.
- ☐ B It converts nitrogen oxides and carbon monoxide to nitrogen and carbon dioxide by being able to change its oxidation state.
- ☐ C It oxidizes unburnt fuel to carbon monoxide.
- ☐ D It oxidizes unburnt fuel to carbon dioxide.

(Total for Question 6 = 1 mark)

7 Which of the following gives the electronic configuration for chromium and for the  $\text{Cr}^{3+}$  ion?

|                          |   | Cr                                  | $\text{Cr}^{3+}$                    |
|--------------------------|---|-------------------------------------|-------------------------------------|
| <input type="checkbox"/> | A | $[\text{Ar}]3\text{d}^44\text{s}^2$ | $[\text{Ar}]3\text{d}^34\text{s}^0$ |
| <input type="checkbox"/> | B | $[\text{Ar}]3\text{d}^54\text{s}^1$ | $[\text{Ar}]3\text{d}^24\text{s}^1$ |
| <input type="checkbox"/> | C | $[\text{Ar}]3\text{d}^54\text{s}^1$ | $[\text{Ar}]3\text{d}^34\text{s}^0$ |
| <input type="checkbox"/> | D | $[\text{Ar}]3\text{d}^44\text{s}^2$ | $[\text{Ar}]3\text{d}^14\text{s}^2$ |

(Total for Question 7 = 1 mark)

8 Which of the following gives the colours of solutions containing  $\text{Cr}_2\text{O}_7^{2-}$ ,  $\text{CrO}_4^{2-}$ , hydrated  $\text{Cr}^{3+}$  and hydrated  $\text{Cr}^{2+}$ , in this order?

- ☐ A Yellow, orange, green, blue
- ☐ B Orange, yellow, green, blue
- ☐ C Orange, yellow, blue, green
- ☐ D Orange, green, yellow, blue

(Total for Question 8 = 1 mark)

9 When dichromate(VI) ions,  $\text{Cr}_2\text{O}_7^{2-}$ , react with iron(II) ions in acidic solution, the products are chromium(III) ions and iron(III) ions. In what ratio do the dichromate(VI) ions and the iron(II) ions react?

- ☐ A 1:6
- ☐ B 1:5
- ☐ C 2:5
- ☐ D 1:3

(Total for Question 9 = 1 mark)

10 The compound  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$  is

- ☐ A tetrahedral with no isomers.
- ☐ B square planar with no isomers.
- ☐ C tetrahedral with two isomers.
- ☐ D square planar with two isomers.

(Total for Question 10 = 1 mark)

11 The hexaaquacopper(II) ion,  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$ , is blue because the water ligands

- ☐ A split the  $p$ -orbital energies and  $p$ - $p$  electron transitions emit blue light.
- ☐ B split the  $d$ -orbital energies and  $d$ - $d$  electron transitions absorb all but blue light.
- ☐ C split the  $p$ -orbital energies and  $p$ - $p$  electron transitions absorb all but blue light.
- ☐ D split the  $d$ -orbital energies and  $d$ - $d$  electron transitions emit blue light.

(Total for Question 11 = 1 mark)

**12** If phenol and benzene are tested separately with bromine water, you would expect to see that

- ☐ **A** benzene and phenol would both decolorize bromine water.
- ☐ **B** benzene would decolorize bromine water, but phenol would not do so.
- ☐ **C** neither benzene nor phenol would decolorize bromine water.
- ☐ **D** benzene would not decolorize bromine water, but phenol would do so.

(Total for Question 12 = 1 mark)

**13** An organic compound **X** is much more soluble in dilute hydrochloric acid than in water. Compound **X** forms a coloured complex with aqueous copper(II) ions.

Compound **X** could be

- ☐ **A**  $\text{C}_6\text{H}_5\text{COOH}$
- ☐ **B**  $\text{C}_6\text{H}_5\text{NO}_2$
- ☐ **C**  $\text{C}_6\text{H}_5\text{NH}_2$
- ☐ **D**  $\text{C}_6\text{H}_5\text{OH}$


(Total for Question 13 = 1 mark)

**14** Which of the following shows the generation of the electrophile in the reaction of benzene with ethanoyl chloride in the presence of anhydrous aluminium chloride?

- ☐ **A**  $\text{CH}_3\text{COCl} + \text{AlCl}_3 \rightarrow [\text{CH}_3\text{CO}]^+ + \text{AlCl}_4^-$
- ☐ **B**  $\text{CH}_3\text{COCl} + \text{AlCl}_3 \rightarrow [\text{CH}_3\text{CO}]^- + \text{AlCl}_4^+$
- ☐ **C**  $\text{CH}_3\text{CH}_2\text{Cl} + \text{AlCl}_3 \rightarrow [\text{CH}_3\text{CH}_2]^+ + \text{AlCl}_4^-$
- ☐ **D**  $\text{CH}_3\text{COOCl} + \text{AlCl}_3 \rightarrow [\text{CH}_3\text{COO}]^- + \text{AlCl}_4^+$

(Total for Question 14 = 1 mark)

15 Which of the following reagents and conditions would enable phenylamine to be converted to the yellow dye 4-hydroxyazobenzene in a good yield?



- ☐ A Sodium nitrite,  $\text{NaNO}_2$ , in concentrated  $\text{HCl}$ , between  $0^\circ\text{C}$  and  $10^\circ\text{C}$ ; followed by an alkaline solution of phenol.
- ☐ B Sodium nitrite,  $\text{NaNO}_2$ , in concentrated  $\text{HCl}$ , between  $0^\circ\text{C}$  and  $10^\circ\text{C}$ ; followed by an acidic solution of phenol.
- ☐ C Sodium nitrate,  $\text{NaNO}_3$ , in concentrated  $\text{HCl}$ , between  $0^\circ\text{C}$  and  $10^\circ\text{C}$ ; followed by an alkaline solution of phenol.
- ☐ D Sodium nitrite,  $\text{NaNO}_2$ , in concentrated  $\text{HCl}$ , room temperature; followed by an alkaline solution of phenol.

(Total for Question 15 = 1 mark)





16 1-butylamine,  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$ , reacts with ethanoyl chloride to form

- ☐ A  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_3^+\text{Cl}^-$
- ☐ B  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{NHCOCH}_3$
- ☐ C  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{NHCH}_2\text{CH}_3$
- ☐ D  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{COCH}_3)\text{NH}_2$

(Total for Question 16 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.


17 The addition polymer poly(ethenol) is water-soluble. The repeating unit of poly(ethenol) is




(Total for Question 17 = 1 mark)

18 A white organic compound, **X**, is optically active and reacts with ninhydrin to give a coloured product. The structural formula of **X** could be


☐ **A**




☐ **B**



☐ **C**



☐ **D**



(Total for Question 18 = 1 mark)

19 Glycine,  $\text{H}_2\text{NCH}_2\text{COOH}$ , is a solid that has a melting temperature of about  $250^\circ\text{C}$ , and it is very soluble in water. This is because of the

- ☐ A formation of intermolecular hydrogen bonds in the solid and hydrogen bonds with water.
- ☐ B formation of  $\text{H}_3\text{N}^+\text{CH}_2\text{COO}^-$  ions which interact strongly with each other in the solid and with water.
- ☐ C dissociation of the molecule to form  $\text{H}_2\text{NCH}_2\text{COO}^-$  and  $\text{H}^+$  ions in the solid and the solution.
- ☐ D protonation of the molecule to form  $\text{H}_3\text{N}^+\text{CH}_2\text{COOH}$  ions in both the solid and the solution.

(Total for Question 19 = 1 mark)

20 Organic solids are often purified by recrystallization. This technique works on the basis that

- ☐ A the impurities must be insoluble in the solvent used.
- ☐ B the impurities must react with the solvent used.
- ☐ C the impurities crystallize first when the hot solution is cooled.
- ☐ D the cooled solution is saturated with the desired material but not with the impurities.

(Total for Question 20 = 1 mark)

**TOTAL FOR SECTION A = 20 MARKS**


## SECTION B

**21** Brass is an alloy of copper, zinc and, in some cases, other metals too. There are over 30 varieties of brass for different applications.

The amount of copper in a brass can be found as follows:

- A weighed sample of brass is reacted with the minimum amount of concentrated nitric acid.
- The solution is neutralized, a portion of it pipetted into a conical flask, and excess potassium iodide solution is added.
- The iodine produced is titrated with a solution of sodium thiosulfate of known concentration.

(a) The ionic equation for the reaction between copper metal and concentrated nitric acid is shown below.



- (i) Give the oxidation numbers of the copper and nitrogen in both the reactants and products.

(2)

**Copper:** from ..... to .....

**Nitrogen:** from ..... to .....

- (ii) Write the two half-equations that can be combined to give the ionic equation shown above.

(2)

- (iii) Explain why the standard electrode potentials for the two ionic half-equations that you have written give an incorrect value for  $E_{\text{cell}}$  for this reaction as described above.

(2)

.....

.....

.....

- (b) The solution produced contains a mixture of zinc ions and copper ions.

- (i) State TWO observations that you would see if concentrated ammonia solution were to be added, drop by drop, to the solution until in excess.

(2)

.....

.....

.....

- (ii) Copper ions can be separated from the zinc ions in the solution by adding sodium hydroxide solution in excess, followed by filtration of the mixture.

Write equations, including state symbols, for the THREE reactions that occur.

(3)

**Equation 1**

**Equation 2**

**Equation 3**

\*(iii) Give examples of amphoteric behaviour and ligand exchange, by reference to the reactions of zinc compounds.

(3)

---

---

---

---

---

---


---

(c) A sample of Admiralty Brass of mass 3.00 g was treated with nitric acid and made up to a neutral solution of volume 250 cm<sup>3</sup>. Excess potassium iodide was added to 25.0 cm<sup>3</sup> portions of this solution, and the liberated iodine was titrated with sodium thiosulfate solution, concentration 0.100 mol dm<sup>-3</sup>. The mean titre was 33.10 cm<sup>3</sup>.

(i) Write the ionic equation for the reaction between thiosulfate ions and iodine.

(1)

\*(ii) The equation for the reaction between copper(II) ions and iodide ions is shown below.



Hence calculate the percentage by mass of copper in Admiralty Brass. Give your answer to **three** significant figures.

(6)

- (iii) When setting up the burette, a student failed to fill the jet of the burette.  
Explain the effect that this would have on the value of the first titre.

If this first titre was included in the calculation of the mean titre, what effect  
would this have on the value for the percentage of copper in the brass?


(2)

---

(Total for Question 21 = 23 marks)

---

22 Paracetamol is possibly the most widely used analgesic (painkiller) in the world. It can be made from phenol as shown below.



- (a) The nitration of phenol in **step 1** uses dilute nitric acid at room temperature, whereas the nitration of benzene requires a mixture of concentrated nitric and sulfuric acids at about 55°C.
- (i) Give the mechanism for the nitration of **benzene**, including the equation for the reaction that produces the electrophile.

(4)

\*(ii) Explain why phenol can be nitrated under much milder conditions than those required to nitrate benzene.

(2)


(iii) Suggest reagents that could be used for **step 2**.

(2)


(iv) Suggest the name or formula of a reagent that could be used in **step 3**.

(1)

\*(b) In the nitration of phenol, **step 1**, two compounds are produced.



**2-nitrophenol**



**4-nitrophenol**

These compounds can be separated by steam distillation, since 2-nitrophenol is volatile in steam but 4-nitrophenol is not.

Describe briefly the technique of steam distillation and give ONE advantage of steam distillation over normal distillation.


(3)

- (c) The nmr spectrum of paracetamol is given below. The peaks at 6.7 and 7.4 ppm are due to the protons on the benzene ring and are both doublets.

Explain why these two peaks are doublets but all the other peaks are singlets.

[There is no need to refer to your data booklet]

(2)



(Total for Question 22 = 14 marks)

**23** More than half of the elements in the Periodic Table are transition elements. Vanadium, element 23, is a typical transition element.

- (a) (i) Give TWO properties shown by vanadium **compounds** that are characteristic of transition metal chemistry, other than variable oxidation state.

(2)

---

---

- (ii) Vanadium(III) ions in aqueous solution exist as  $[\text{V}(\text{H}_2\text{O})_6]^{3+}$ .

Draw this ion so as to clearly show its shape. Name the type of bond between the ligand and the vanadium ion and state the feature of the ligand that enables this bond to be formed.

(3)

---

---

---


---

(b) Vanadium ions exist in oxidation states from (V) to (II).

- (i) Use your data booklet (page 15) to find the standard electrode (reduction) potential for the reduction of vanadium(IV),  $\text{VO}^{2+}$ , to vanadium(III),  $\text{V}^{3+}$ . (1)

- \* (ii) Explain the term **disproportionation**. (2)

- (iii) Use your answer to (b)(i), and the data below, to calculate  $E^\ominus_{\text{cell}}$  for the formation of vanadium(V) and vanadium(III) from vanadium(IV) in acidic solution. State if the reaction is feasible under standard conditions and justify your answer.



(Total for Question 23 = 10 marks)

TOTAL FOR SECTION B = 47 MARKS

## SECTION C


24

A major function of the chemical industry is the manufacture of perfumes and flavourings. Perfumes were originally made from natural products, such as spermaceti from whales or civetone from the civet cat. The use of synthetic equivalents is generally cheaper and the supply is more reliable, and does not require the animals to be killed which obtaining spermaceti did. Synthetic perfumes and flavourings are usually nature-identical, although naturally-occurring molecules that have stereoisomers are sometimes produced synthetically as mixtures since the stereospecific synthesis required might be difficult to achieve economically.

Some chiral molecules have a different taste depending on which enantiomer is present. The enantiomer (–)-carvone tastes and smells of spearmint, and its mirror-image, (+)-carvone, of caraway or dill. In some cases only one enantiomer has any taste; this is true for glucose. The mirror image molecule of naturally occurring glucose has no taste and cannot even be absorbed or metabolized. Many drug molecules are chiral, though paracetamol is not. The wrong isomer present in a drug may be positively damaging, which was the case with thalidomide. In order to synthesize optically-pure drug molecules, it is important to understand the mechanism of any reaction used. Using an  $S_N1$  reaction which involves the chiral centre would result in the product mixture being racemic. It is advantageous to use stereospecific catalysts wherever possible, and industry on the whole prefers to use heterogeneous rather than homogeneous catalysts.

- (a) The skeletal formula of carvone is shown below. Draw a circle around the chiral carbon atom.

(1)



(b) Explain why the synthesis of paracetamol is more efficient than the synthesis of a single enantiomer such as (–)-carvone.

(3)

---

---

---

---

---

(c) Carvone contains two types of functional group. For each of these, give a suitable chemical test to show its presence, and state what you would see in each case.

(4)

---

---

---

---

(d) Carvone can be reduced in a variety of ways.

- (i) On reduction with hydrogen, in the presence of a platinum catalyst, 4.5 g of carvone reacted with 1.44 dm<sup>3</sup> of hydrogen. Use these data to deduce the skeletal formula of the reduction product.


[Molar mass of carvone is 150 g mol<sup>-1</sup>; molar volume of hydrogen at the temperature and pressure of the experiment is 24 dm<sup>3</sup> mol<sup>-1</sup>.]

(3)

**Working**

**Hence skeletal formula of reduction product.**

- (ii) Reduction of carvone, using hydrazine in potassium hydroxide solution, gives limonene.



How would you show from a comparison of the infra-red spectra of carvone and of limonene that this reduction had occurred? You should quote appropriate data, from the data booklet, to support your answer.

(2)

.....

.....

.....

.....

.....

- (e) (i) Draw the skeletal formula of the molecule that would be obtained if carvone were to be reacted with an excess of hydrogen bromide in an inert solvent.

(2)

- (ii) If the product from (e)(i) were to be heated with ethanolic potassium hydroxide solution, elimination would occur and HBr would be lost. Would the resulting molecule necessarily be carvone? Explain your answer.

(2)

.....

.....

.....

.....

- (f) \*(i) Give the mechanism of one nucleophilic reaction of your choice, which **either** gives rise to a racemic mixture **or** results in the inversion of the chirality of the starting material. Explain what your mechanism predicts about the stereochemistry of the product.

(5)

- (ii) Suggest why industrial chemists prefer to use heterogeneous rather than homogeneous catalysts.

(1)

(Total for Question 24 = 23 marks)

**TOTAL FOR SECTION C = 23 MARKS**  
**TOTAL FOR PAPER = 90 MARKS**

mock papers 2

SECTION A

1 What type of bonding occurs between the metal ion and ligand in the complex ion  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$ ?

- ☐ A Metallic
- ☐ B Ionic
- ☐ C Hydrogen
- ☐ D Dative covalent

(Total for Question 1 = 1 mark)

2 Which of these four amino acids could **not** rotate the plane of plane-polarised light?

- ☐ A  $\text{H}_2\text{NCH}(\text{CH}_3)\text{COOH}$
- ☐ B  $\text{H}_2\text{NCH}(\text{CH}_2\text{COOH})\text{COOH}$
- ☐ C  $\text{H}_2\text{NCH}_2\text{COOH}$
- ☐ D  $\text{H}_2\text{NCH}(\text{CH}_2\text{SH})\text{COOH}$

(Total for Question 2 = 1 mark)

3 In the solid state, the amino acid serine exists in the form

- ☐ A  $\text{H}_3\text{N}^+\text{CH}(\text{CH}_2\text{OH})\text{COOH}$
- ☐ B  $\text{H}_3\text{N}^+\text{CH}(\text{CH}_2\text{OH})\text{CO}_2^-$
- ☐ C  $\text{H}_2\text{NCH}(\text{CH}_2\text{OH})\text{COOH}$
- ☐ D  $\text{H}_2\text{NCH}(\text{CH}_2\text{OH})\text{CO}_2^-$

(Total for Question 3 = 1 mark)

4 The best method for separating a mixture of amino acids in solution is

- ☐ A distillation.
- ☐ B solvent extraction.
- ☐ C chromatography.
- ☐ D recrystallization.

(Total for Question 4 = 1 mark)

5 How many different peaks due to hydrogen atoms would you expect to see in a **low resolution** proton nmr spectrum of propanoic acid,  $\text{CH}_3\text{CH}_2\text{COOH}$ ?

- ☐ A Two
- ☐ B Three
- ☐ C Five
- ☐ D Six

(Total for Question 5 = 1 mark)

6 In a **high resolution** proton nmr spectrum of ethanoic acid,  $\text{CH}_3\text{COOH}$ , the peak due to the hydrogen atoms in the methyl group would be a

- ☐ A singlet.
- ☐ B doublet.
- ☐ C triplet.
- ☐ D quartet.

(Total for Question 6 = 1 mark)



7 Which of these compounds will **not** form an amide in a reaction with ethanoyl chloride?

- ☐ A  $\text{NH}_3$
- ☐ B  $\text{CH}_3\text{CH}_2\text{NH}_2$
- ☐ C  $\text{CH}_3\text{CH}_2\text{NH}(\text{CH}_3)$
- ☐ D  $\text{CH}_3\text{CH}_2\text{N}(\text{CH}_3)_2$

(Total for Question 7 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.

8 This question concerns the following organic compounds.



Which compound is most likely to

(a) form the solution with the lowest pH when mixed with water?

(1)

☐ A

☐ B

☐ C

☐ D

(b) burn with a smoky flame?

(1)

☐ A

☐ B

☐ C

☐ D

(c) have a fruity smell?

(1)

☐ A

☐ B

☐ C

☐ D

(d) have an absorption in its IR spectrum at about  $1795\text{ cm}^{-1}$ ?

(1)

☐ A


☐ B

☐ C

☐ D

(Total for Question 8 = 4 marks)

9 This question is about the reaction scheme below.



Which step is most likely to need

(a) tin and concentrated hydrochloric acid?

(1)

- ☐ A Step 1
- ☐ B Step 2
- ☐ C Step 3
- ☐ D Step 4

(b) a catalyst of iron(III) chloride?

(1)

- ☐ A Step 1
- ☐ B Step 2
- ☐ C Step 3
- ☐ D Step 4



(c) a nickel catalyst?

(1)

- ☐ A Step 1
- ☐ B Step 2
- ☐ C Step 3
- ☐ D Step 4

(Total for Question 9 = 3 marks)

10 The apparatus below can be used to measure the value of  $E_{\text{cell}}$  for the reaction




The electrodes are:

|                          |   | electrode 1 | electrode 2 |
|--------------------------|---|-------------|-------------|
| <input type="checkbox"/> | A | zinc        | iron        |
| <input type="checkbox"/> | B | iron        | zinc        |
| <input type="checkbox"/> | C | zinc        | platinum    |
| <input type="checkbox"/> | D | platinum    | platinum    |

(Total for Question 10 = 1 mark)

11 Copper reacts with silver ions according to the reaction below.



$E_{\text{cell}}^{\ominus}$  for this reaction is

- ☐ A +0.46 V
- ☐ B +1.14 V
- ☐ C +1.26 V
- ☐ D +1.94 V

(Total for Question 11 = 1 mark)

12  $E_{\text{cell}}^{\ominus}$  for four reactions are shown in the table below.

|            | $E_{\text{cell}}^{\ominus} / \text{V}$ |
|------------|----------------------------------------|
| Reaction 1 | +1.10                                  |
| Reaction 2 | +0.65                                  |
| Reaction 3 | +0.10                                  |
| Reaction 4 | -1.30                                  |

Which reaction

(a) is thermodynamically not feasible?

(1)

- ☐ A Reaction 1
- ☐ B Reaction 2
- ☐ C Reaction 3
- ☐ D Reaction 4


(b) has the largest value for  $\ln K$ ?

(1)

- ☐ A Reaction 1
- ☐ B Reaction 2
- ☐ C Reaction 3
- ☐ D Reaction 4

(Total for Question 12 = 2 marks)


13 Consider the reaction scheme below and calculate the mass of aspirin you would expect to form if you started with 47 g of phenol.



- ☐ A 31.96 g
- ☐ B 61.20 g
- ☐ C 74.25 g
- ☐ D 90.00 g

(Total for Question 13 = 1 mark)

14 Which of the monomers **A to D** would form the polymer below?




- ☐ A
- ☐ B
- ☐ C
- ☐ D

(Total for Question 14 = 1 mark)

TOTAL FOR SECTION A = 20 MARKS

SECTION B

- 15 In the reaction shown below, the aromatic compound 1,4-dimethylbenzene reacts with 2-bromobutane. The reaction is catalysed by aluminium chloride,  $\text{AlCl}_3$ , which dissolves in the reaction mixture.



- (a) (i) Name the type of reaction and the mechanism.

(1)

- (ii) Write the equation to show how the attacking species forms and give the mechanism for the reaction.

(4)


**Equation**

**Mechanism**

- (b) The same reaction can also be carried out using a heterogeneous graphite catalyst under similar conditions. Assuming both reactions have a similar rate and yield, suggest one advantage of using the solid graphite catalyst instead of aluminium chloride. Justify your answer.

(2)

- (c) A student proposed to synthesise the compound 2,5-dimethylphenylamine, used in the manufacture of dyes, following the scheme below.



- (i) What two reagents are needed for **step 1**?

(2)

- (ii) Suggest why 1,4-dimethylbenzene is more reactive than benzene in reactions such as **step 1**.

(2)

(1)

(5)

(Total for Question 15 = 17 marks)

**16** The leaves of the rhubarb plant contain ethanedioic acid,  $(\text{COOH})_2$ , a toxic white soluble solid. The acid is readily oxidized by potassium manganate(VII) under acidic conditions. A sample of 250 g of rhubarb leaves was finely chopped then soaked in warm water to release any ethanedioic acid present. The mixture was then filtered and made up to a volume of  $500 \text{ cm}^3$  using distilled water.  $10.0 \text{ cm}^3$  of the solution was then titrated with  $0.0100 \text{ mol dm}^{-3}$  acidified potassium manganate(VII) solution from a burette, requiring  $11.30 \text{ cm}^3$  to completely oxidize the sample.

- (a) (i) Write the half equation for the oxidation of ethanedioic acid to form carbon dioxide, and the half equation for the reduction of manganate(VII) ions,  $\text{MnO}_4^-$ , in acidic solution to form manganese(II) ions. State symbols are **not** required. (2)

- (ii) Use your answers to (a)(i) to write the overall equation for the reaction, showing that the ratio of ethanedioic acid to manganate(VII) ions in the full equation is 5 : 2. State symbols are **not** required. (1)

- \*(iii) Calculate the % by mass of the ethanedioic acid present in the leaves, giving your final answer to **two** decimal places.

(5)

- (iv) What is the level of accuracy of a burette in each reading? Use your answer to calculate the percentage error in the titre volume of  $11.30 \text{ cm}^3$ .

(2)

- (v) Suggest **two** reasons, other than the accuracy of the equipment used for measurements, why the results obtained in this experiment may be considered unreliable.

(2)

- (vi) A student risk assessment for this experiment suggested wearing gloves, but a supervisor said that this was unnecessary. Why do you think this precaution was suggested by the student and why was it rejected by the supervisor?

(2)

- (vii) An aqueous solution of  $\text{MnO}_4^-$  ions contained a small amount of chloride ions,  $\text{Cl}^-$ , as an impurity. Use this fact, and items 70 and 85 from page 16 of the data booklet, to suggest why this solution went cloudy after a time.

(2)

(b) An aqueous solution containing  $\text{Mn}^{2+}$  ions is pale pink in colour due to the presence of the complex ion  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}(\text{aq})$ .

(i) Complete the electronic configuration of the  $\text{Mn}^{2+}$  ion.

(1)

$1s^2$  .....


(ii) What shape would you expect this complex ion to be?

(1)

.....

**(Total for Question 16 = 18 marks)**

17 The reaction scheme below summarises some of the reactions of chromium ions in aqueous solution. Look carefully at the scheme and answer the questions that follow.



(a) (i) Explain why the  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  ion is coloured.

(3)

.....

.....

.....

.....

.....

.....

(ii) Suggest what reagent is needed for **reaction 1** and identify the type of reaction.


(2)

.....

.....

- (b) (i)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  ions react with water to form an acidic solution. Complete the equation for this reaction.

(2)



- (ii) The pH of an aqueous solution of  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$  is higher than that of an aqueous solution of  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  of the same concentration. Suggest why this is so.

(2)

.....

.....

.....

- (c) Give the formula of the green precipitate formed in **reactions 2 and 3**.

(1)

.....

- (d) By considering the nature of the reactants in **reaction 4**, explain why the green precipitate reacts as shown in the scheme. Suggest how you could reverse **reaction 4**.

(3)

.....

.....

.....

.....

.....

.....

- (e) Write the equation for **reaction 6** and use this to explain, in terms of the entropy change, why the complex  $[\text{Cr}(\text{edta})]^-$  is relatively more stable than  $[\text{Cr}(\text{NH}_3)_6]^{3+}$ .

(2)

**Equation**

.....

.....

.....


.....

(Total for Question 17 = 15 marks)


**TOTAL FOR SECTION B = 50 MARKS**

## SECTION C

- 18 Eugenol, a pale yellow oil, and eugenol ethanoate are phenol-derived compounds found in the evergreen clove tree *Eugenia aromaticum*.



Eugenol




Eugenol ethanoate

Eugenol is used in perfumes, the manufacture of food flavourings and as a local anaesthetic. Eugenol ethanoate is mainly used in perfumes and aftershaves. Although used for many years, both compounds are classified as harmful and have been tested to determine their toxicity by ingestion. However, humans would need to consume very large amounts to reach toxic levels.

The compounds are the main constituents of clove oil which can be extracted from the dried buds of *Eugenia aromaticum*'s flowers. Traditionally the oil is extracted by steam distillation, though a greater yield of oil can be obtained using a Soxhlet extractor to pass a chlorinated solvent through the dried buds several times to dissolve the clove oil. An alternative technique uses carbon dioxide as a solvent. Above a temperature of 304 K and a high pressure of 73.8 atm, carbon dioxide behaves as a supercritical fluid and when passed through the clove buds, it dissolves the clove oil. Releasing the pressure causes the carbon dioxide to turn back into a gas, leaving the clove oil behind. A summary of the characteristics of the clove oil obtained by the three extraction techniques is shown in the table below.

| Extraction method            | Mass of oil per 100 g of dried buds / g | % eugenol and eugenol ethanoate in the oil produced | Extraction time / h | Colour and texture | Use of organic solvent |
|------------------------------|-----------------------------------------|-----------------------------------------------------|---------------------|--------------------|------------------------|
| Supercritical carbon dioxide | 19.6                                    | 78.4                                                | 2                   | pale yellow oil    | no                     |
| Steam distillation           | 11.5                                    | 53.5                                                | 4–6                 | brown-yellow oil   | yes                    |
| Soxhlet extraction           | 41.8                                    | 40.1                                                | 6                   | brown paste        | yes                    |

Both molecules can also be manufactured synthetically in the laboratory. A reaction scheme for synthesising both molecules is summarised below.



- (a) (i) 0.328 g of eugenol produced synthetically was burnt completely in excess oxygen, producing 0.880 g of carbon dioxide and 0.216 g of water. Use these data to show they are consistent with the molecular formula of eugenol.

(4)

- (ii) Describe a chemical test you could carry out to confirm the presence of a carbon-carbon double bond in the product of **step 1**. What would you expect to see?

(2)

- (iii) What technique would you use to heat the reactants in **step 3** to minimise the loss of any volatile material?

(1)

- (iv) Suggest what reagent(s) could be used in **step 4**.

(1)

- (b) (i) Draw and label the apparatus suitable for extracting clove oil from clove buds by steam distillation in the laboratory.

(3)

- (ii) The distillate formed is a mixture of water and clove oil with a significant amount of oil dissolved in the water. Outline the steps that have to be taken to obtain the dry oil.

(3)

.....

.....

.....

.....

.....

.....

- (c) Toxicity data for substances such as eugenol are generally obtained by tests on animals such as rats and guinea pigs. In the case of eugenol, do you think such tests are reasonable? Briefly justify your answer.

(1)

- \*(d) Evaluate the three extraction methods for obtaining clove oil using information from the table. Give **one** reason why the synthetic route of obtaining eugenol, shown on page 20, is less preferable than extraction from clove buds.

(5)




(Total for Question 18 = 20 marks)

**TOTAL FOR SECTION C = 20 MARKS**  
**TOTAL FOR PAPER = 90 MARKS**

mock papers 3

SECTION A

1 Four complex ions have the following formulae:



(a) Which complex ion is most likely to be tetrahedral in shape?

(1)

☐ A

☐ B

☐ C

☐ D

(b) Which complex ion is most likely **not** to be coloured?

(1)

☐ A

☐ B

☐ C

☐ D

(c) Each of these complex ions may be formed by ligand exchange from an aqua complex. For which complex ion is the entropy change of this reaction most positive?

(1)

☐ A

☐ B

☐ C

☐ D

(Total for Question 1 = 3 marks)

2 When a **few drops** of aqueous ammonia are added to a solution containing  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  ions the product formed will be

- ☐ A  $[\text{Cr}(\text{NH}_3)_6]^{3+}$
- ☐ B  $\text{Cr}(\text{H}_2\text{O})_3(\text{OH})_3$
- ☐ C  $[\text{Cr}(\text{NH}_3)_4]^{3+}$
- ☐ D  $[\text{Cr}(\text{H}_2\text{O})_2(\text{OH})_4]^-$

(Total for Question 2 = 1 mark)





3 Which of these statements about a standard hydrogen electrode, for which  $E^\ominus = 0 \text{ V}$ , is **not** correct?

- ☐ A The hydrogen gas is at a pressure of 1 atm.
- ☐ B A solution containing  $1 \text{ mol dm}^{-3}$  of  $\text{H}^+(\text{aq})$  ions is used.
- ☐ C A platinum electrode is used.
- ☐ D The temperature is kept at  $20^\circ\text{C}$ .

(Total for Question 3 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.

4 Four polymers labelled **A** to **D** have the following formulae:



(a) Which polymer is most soluble in hot water?

(1)

☐ **A**


☐ **B**

☐ **C**

☐ **D**

(b) Which polymer is formed from the monomer shown below?

(1)



☐ A

☐ B

☐ C

☐ D

(c) Which polymer is a condensation polymer?

(1)

☐ A

☐ B


☐ C

☐ D

(Total for Question 4 = 3 marks)

Use this space for any rough working. Anything you write in this space will gain no credit.

5 The formulae of some organic compounds labelled **A to D** are shown below.



(a) Which compound reacts with sodium to form hydrogen?

(1)

- ☐ A
- ☐ B
- ☐ C
- ☐ D

(b) Which compound forms a green complex ion with  $\text{CuSO}_4(\text{aq})$ ?

(1)

- ☐ A
- ☐ B
- ☐ C
- ☐ D

(c) Which compound forms an orange precipitate with 2,4-dinitrophenylhydrazine?

(1)

- ☐ A
- ☐ B
- ☐ C
- ☐ D

(Total for Question 5 = 3 marks)

6 How many peaks would you expect to see in a **low resolution** proton nmr spectrum of the ester  $\text{HCOOCH}_2\text{CH}_2\text{CH}_3$ ?

- ☐ A 8  
☐ B 7  
☐ C 4  
☐ D 3

(Total for Question 6 = 1 mark)

7 In a **high resolution** proton nmr spectrum of ethyl ethanoate,  $\text{CH}_3\text{COOCH}_2\text{CH}_3$ , the peak due to the hydrogen atoms shown **in bold** would be a

- ☐ A singlet.  
☐ B doublet.  
☐ C triplet.  
☐ D quartet.

(Total for Question 7 = 1 mark)

8 Which of these compounds, whose formulae are shown below, **cannot** exist as a racemic mixture?

- ☐ A  $\text{CH}_2\text{ClCHClCOOH}$   
☐ B  $\text{HOOCCHClCOOH}$   
☐ C  $\text{CH}_3\text{CHClCOOH}$   
☐ D  $\text{CH}_3\text{CH}(\text{OH})\text{COOH}$

(Total for Question 8 = 1 mark)

9 Phenol reacts with bromine water whereas benzene reacts with bromine in the presence of iron.

(a) The mechanism for both these reactions is

(1)

- ☐ A electrophilic substitution.
- ☐ B electrophilic addition.
- ☐ C nucleophilic substitution.
- ☐ D nucleophilic addition.

(b) In the reaction of benzene with bromine, iron

(1)

- ☐ A acts as a heterogeneous catalyst.
- ☐ B acts as a homogeneous catalyst.
- ☐ C reacts with the bromine to make iron(III) bromide, FeBr<sub>3</sub>.
- ☐ D allows bromine to attack the hydrogen atoms on benzene more readily.

(c) Bromine reacts more readily with phenol than with benzene because the OH group on phenol

(1)

- ☐ A is a good leaving group.
- ☐ B attracts the bromine particles more readily.
- ☐ C is a good nucleophile.
- ☐ D increases the electron density of the ring.

(Total for Question 9 = 3 marks)

10 Ammonia (NH<sub>3</sub>), butylamine (CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>) and phenylamine (C<sub>6</sub>H<sub>5</sub>NH<sub>2</sub>) all form alkaline solutions in water. The order of **increasing** pH of equimolar solutions is

- ☐ A C<sub>6</sub>H<sub>5</sub>NH<sub>2</sub> < CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub> < NH<sub>3</sub>
- ☐ B NH<sub>3</sub> < CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub> < C<sub>6</sub>H<sub>5</sub>NH<sub>2</sub>
- ☐ C C<sub>6</sub>H<sub>5</sub>NH<sub>2</sub> < NH<sub>3</sub> < CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>
- ☐ D CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub> < NH<sub>3</sub> < C<sub>6</sub>H<sub>5</sub>NH<sub>2</sub>

(Total for Question 10 = 1 mark)

**11** The distance on a chromatogram moved by an individual amino acid, in a mixture of different amino acids, mainly depends on

- ☐ **A** the molar mass of the amino acid.
- ☐ **B** the molar mass of the solvent used.
- ☐ **C** the intermolecular forces between the solvent and the stationary phase.
- ☐ **D** the intermolecular forces between the amino acid and both the solvent and the stationary phase.

**(Total for Question 11 = 1 mark)**

---

**12** Amino acids are crystalline solids with a high melting temperature because

- ☐ **A** each molecule has a large number of electrons.
- ☐ **B** each molecule forms hydrogen bonds at both ends.
- ☐ **C** a proton is transferred from one end of the molecule to the other.
- ☐ **D** their shape allows the molecules to pack close together.

**(Total for Question 12 = 1 mark)**


---

**TOTAL FOR SECTION A = 20 MARKS**

---

SECTION B

13 Some reactions of benzene are shown below.



(a) (i) Suggest the formula of reagent A in **reaction 1**.

(1)

(ii) Write the equation to show how the catalyst,  $\text{AlCl}_3$ , reacts with reagent A to form the species which attacks the benzene ring.

(1)

(iii) Draw the structure of the intermediate ion formed when the species in (ii) attacks the benzene ring.

(1)

(b) The methylbenzene formed in **reaction 1** generally reacts in a similar way to benzene but faster, as the ring is said to be activated.

(i) Explain how the presence of a methyl group activates the benzene ring.

(1)

(ii) Use your answer to (i) to explain why methylbenzene reacts faster.

(1)

(c) (i) Draw the structural formula of **compound X**, formed in **reaction 2**.

(1)

(ii) The organic product of **reaction 2** is also formed when the same reactants, but with an aluminium catalyst, are heated using microwave radiation. Suggest two reasons why this technique may be considered 'greener'.

(2)


(d) Name reagent B needed for **reaction 3**.

(1)

(Total for Question 13 = 9 marks)

14 This question is about synthetically produced painkillers and anaesthetics.

- (a) The local anaesthetic procaine can be synthesised from benzoic acid. The simplified route is shown below.



- (i) Suggest the two reagents needed for **step 1**.

(2)

(ii) Draw the apparatus needed to heat under reflux in **step 2**.

(3)

(iii) Suggest why the reagents for the reaction in **step 2** are

(2)

**heated** .....

.....

**under reflux** .....

.....

(iv) Give the structural formula for the organic reagent needed in **step 3**.

(1)

(v) What type of reaction is taking place in **step 4**? Suggest the reagents used.

(2)


.....

.....

.....

.....

- (b) A student produced a sample of aspirin by the esterification of 9.40 g of 2-hydroxybenzoic acid with excess ethanoic anhydride.



After purification by recrystallization, 7.77 g of aspirin was obtained.

[ $M_r$  of 2-hydroxybenzoic acid = 138,  $M_r$  of aspirin = 180]

- (i) Calculate the percentage yield obtained.

(3)

- \*(ii) Outline how to purify a solid, such as aspirin, by recrystallization, using water as the solvent.

(4)

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

(iii) Explain what effect recrystallization has on the final yield.

(1)

(c) Paracetamol is found in many non-prescription painkillers, often in conjunction with other compounds such as codeine.



paracetamol

(i) Suggest, by name or formula, a reagent that could be used to form paracetamol from 4-aminophenol.

(1)

(ii) Suggest why sales of non-prescription painkillers, often containing paracetamol and codeine, are limited to 32 tablets.

(1)

(iii) Explain why paracetamol is only slightly soluble in water although it can form hydrogen bonds with water.

(1)

(Total for Question 14 = 21 marks)

**15** Hydrogen gas can be used as a fuel in car engines by being burnt in a combustion reaction or reacted with oxygen in a fuel cell to produce electricity.

- (a) Write half-equations for the reaction of hydrogen gas at the anode and oxygen gas at the cathode in the fuel cell.

(2)

Anode

Cathode

- (b) Describe one advantage of using hydrogen in fuel cells rather than burning the hydrogen directly.

(1)

.....

.....

.....

.....

- (c) Other fuels, such as ethanol, can also be used in fuel cells. By considering the possible sources of ethanol and hydrogen, explain why some scientists believe the use of such cells could provide a more sustainable source of energy for cars, compared with fossil fuels.

(3)

.....

.....

.....

.....

.....

.....

**(Total for Question 15 = 6 marks)**

**16** This question is about the transition metal iron and some of its compounds.

- (a) Give the electronic configuration of the  $\text{Fe}^{3+}$  ion and use this to define what is meant by a transition element.

(2)

.....

.....

.....

.....

- \*(b) Iron will act as a surface catalyst in some gaseous reactions. Outline the processes that take place during such catalysis and suggest two reasons to explain why the catalyst speeds up the reaction.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) One of the components of rust, found on objects made from iron, is iron(III) hydroxide,  $\text{Fe}(\text{OH})_3$ . Use items 17, 19 and 44 from the Standard Electrode Potential table in your data booklet to show how it is able to form in two steps, writing an equation for each step.

(4)

- (d) Haemoglobin is a complex containing iron(II) ions.

Describe how nitrogen atoms in the haemoglobin bond to the iron(II) ions.

(2)

.....

.....

.....

.....

---

(Total for Question 16 = 12 marks)

---

**TOTAL FOR SECTION B = 48 MARKS**


## SECTION C

17

Alcoholic drinks contain ethanol,  $\text{CH}_3\text{CH}_2\text{OH}$ , in aqueous solution. The percentage of alcohol in a drink can be determined by a redox titration, whilst the amount of alcohol present on the breath of someone who has consumed such a drink can be estimated using a breathalyser.

The earliest breathalysers used the colour change that occurs when dichromate(VI) ions react with ethanol to measure the amount of alcohol. Later models measure the current from a fuel cell. Cheaper versions of these meters are available for drivers to buy for self-testing. Some police forces also use fuel cell breathalysers in conjunction with infrared breath analysers, which can determine the amounts of alcohol from an infrared spectrum.

In an experiment to find out the concentration of ethanol in a drink, a small beaker containing  $5.00 \text{ cm}^3$  of a diluted sample of the drink is suspended above  $10.0 \text{ cm}^3$  of excess acidified sodium dichromate(VI) solution, of concentration  $0.0800 \text{ mol dm}^{-3}$ , and left for 24 hours in a warm place.



The ethanol vaporizes and reacts with some of the acidified sodium dichromate(VI) ions. Excess potassium iodide is then added to the unreacted acidified sodium dichromate(VI), forming iodine,  $\text{I}_2(\text{aq})$ .

The  $\text{I}_2(\text{aq})$  is then titrated with a solution of sodium thiosulfate,  $\text{Na}_2\text{S}_2\text{O}_3$ , of concentration  $0.0250 \text{ mol dm}^{-3}$ .

(a) Ethanol and dichromate(VI) ions in acidic solution react in the mole ratio 3:2.

(i) Complete the two half-equations below. State symbols are **not** required.

(2)





- (ii) Use either the half-equations in (i) or that the mole ratio of  $\text{CH}_3\text{CH}_2\text{OH}:\text{Cr}_2\text{O}_7^{2-}$  is 3:2 to construct the ionic equation for the reaction between ethanol and acidified dichromate(VI) ions. State symbols are **not** required.

(1)

- \*(iii) The iodine formed in the experiment reacted completely with  $34.40\text{ cm}^3$  of the  $0.0250\text{ mol dm}^{-3}$  sodium thiosulfate solution. Use this information, the fact that the mole ratio of  $\text{CH}_3\text{CH}_2\text{OH}:\text{Cr}_2\text{O}_7^{2-}$  is 3:2 and the equations below, to calculate the concentration of the ethanol in the  $5.00\text{ cm}^3$  of the diluted sample of the drink.

(6)



- (iv) The solution used in the experiment was made by adding  $10.0 \text{ cm}^3$  of the drink to a  $100 \text{ cm}^3$  volumetric flask, making up the remainder of the volume using distilled water. Use the information and your final answer to (a)(iii) to calculate the concentration, in  $\text{mol dm}^{-3}$ , of ethanol in the undiluted drink.

(1)

- (v) Suggest why the sample was suspended above the acidified sodium dichromate(VI) solution, rather than simply being mixed with it.

(1)

- (vi) Suggest two reasons why the apparatus was left in a **warm** place for **24 hours**.  
What would be the effect on the final result if this procedure were not followed?

(3)

- (vii) Do you think that this experiment gives a reliable result? Explain your answer.

(1)

- \* (b) (i) Explain how each type of breathalyser, mentioned in the passage, shows the amount of ethanol present.

(3)

Earliest type.....

.....

.....

Fuel cell.....

.....

.....

Infrared.....

.....

.....

- (ii) Suggest why infrared breathalysers do not use the OH absorption to detect the amount of alcohol on the breath.

(1)

.....

.....

.....

(iii) Suggest why some police forces use infrared breathalysers together with fuel cell breathalysers.

(1)

---

---

---

---

(iv) Suggest **one** advantage and **one** disadvantage of buying a personal breathalyser.

(2)

---

---

---

---

---

---

---

---

---

---

---

(Total for Question 17 = 22 marks)

---

**TOTAL FOR SECTION C = 22 MARKS**

**TOTAL FOR PAPER = 90 MARKS**

mock papers 4



SECTION A

- 1 An electrochemical cell consists of a standard hydrogen electrode and a  $\text{Cu}^{2+}(\text{aq})|\text{Cu}(\text{s})$  electrode which uses copper(II) sulfate solution. Which one of the following does **not** affect the e.m.f. of the cell?


- ☐ A The volume of the copper(II) sulfate solution.
- ☐ B The temperature.
- ☐ C The pressure of the hydrogen.
- ☐ D The concentration of the copper(II) sulfate solution.

(Total for Question 1 = 1 mark)

- 2 Which answer corresponds to the correct value of  $E_{\text{cell}}^{\ominus}$  for the oxidation of hydrogen peroxide by manganate(VII) ions? The half-reactions are



The overall equation is



- ☐ A  $E_{\text{cell}}^{\ominus} = + 2.19 \text{ V}$
- ☐ B  $E_{\text{cell}}^{\ominus} = - 0.83 \text{ V}$
- ☐ C  $E_{\text{cell}}^{\ominus} = - 0.38 \text{ V}$
- ☐ D  $E_{\text{cell}}^{\ominus} = + 0.83 \text{ V}$

(Total for Question 2 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.

3 The transition metal complex  $\text{Pt}(\text{NH}_3)_2\text{Cl}_2$  exists as two geometric isomers. This is because the complex

- ☐ A is square-planar.
- ☐ B is tetrahedral.
- ☐ C contains a double bond.
- ☐ D is octahedral.

(Total for Question 3 = 1 mark)

4 Hydrogen peroxide,  $\text{H}_2\text{O}_2$ , can be analysed by titration. The hydrogen peroxide solution is treated with acidified potassium iodide solution, and the liberated iodine is titrated with a standard solution of sodium thiosulfate,  $\text{Na}_2\text{S}_2\text{O}_3$ . The products are iodide ions and tetrathionate ions,  $\text{S}_4\text{O}_6^{2-}$ .

Which of the following applies to this reaction?

|                          |   | Action of $\text{H}_2\text{O}_2$ | Action of $\text{S}_2\text{O}_3^{2-}$ |
|--------------------------|---|----------------------------------|---------------------------------------|
| <input type="checkbox"/> | A | oxidizing agent                  | oxidizing agent                       |
| <input type="checkbox"/> | B | oxidizing agent                  | reducing agent                        |
| <input type="checkbox"/> | C | reducing agent                   | oxidizing agent                       |
| <input type="checkbox"/> | D | reducing agent                   | reducing agent                        |

(Total for Question 4 = 1 mark)

5 A hydrated transition metal ion is colourless. Which of the following could be the electronic configuration of this ion?

- ☐ A  $[\text{Ar}] 3d^5 4s^2$
- ☐ B  $[\text{Ar}] 3d^8$
- ☐ C  $[\text{Ar}] 3d^{10} 4s^2$
- ☐ D  $[\text{Ar}] 3d^{10}$

(Total for Question 5 = 1 mark)

6 Which of the following reagents would enable you to separate iron(III) hydroxide from a mixture of iron(III) hydroxide and copper(II) hydroxide?

- ☐ A Dilute hydrochloric acid
- ☐ B Aqueous ammonia
- ☐ C Dilute nitric acid
- ☐ D Sodium hydroxide solution

(Total for Question 6 = 1 mark)

7 When a solution containing 0.10 mol of chromium(III) chloride,  $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$ , is treated with excess silver nitrate solution, 0.20 mol of silver chloride,  $\text{AgCl}$ , is immediately precipitated. The formula of the complex ion in the solution is

- ☐ A  $[\text{Cr}(\text{OH})_6]^{3-}$
- ☐ B  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$
- ☐ C  $[\text{CrCl}(\text{H}_2\text{O})_5]^{2+}$
- ☐ D  $[\text{CrCl}_2(\text{H}_2\text{O})_4]^+$

(Total for Question 7 = 1 mark)

8 Which of the following species is **not** able to act as a ligand in the formation of transition metal complexes?

- ☐ A  $\text{C}_6\text{H}_5\text{NH}_2$
- ☐ B  $\text{NH}_3$
- ☐ C  $\text{NH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$
- ☐ D  $\text{NH}_4^+$

(Total for Question 8 = 1 mark)

9 The element zinc, with electronic configuration  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2$ , is **not** regarded as a transition element because

- ☐ A the oxide of zinc is amphoteric.
- ☐ B none of its ions has an unpaired electron in the *d*-subshell.
- ☐ C it does not readily form complex ions.
- ☐ D it has a boiling temperature low enough for it to be easily distilled.





(Total for Question 9 = 1 mark)

10 The compounds below were heated with aqueous sodium hydroxide solution. Which one of them did **not** give sodium ethanoate,  $\text{CH}_3\text{COONa}$ , as one of the products?

- ☐ A  $\text{CH}_3\text{COOCH}_3$
- ☐ B  $\text{CH}_3\text{COCH}_3$
- ☐ C  $\text{CH}_3\text{COOH}$
- ☐ D  $\text{CH}_3\text{COCl}$

(Total for Question 10 = 1 mark)

11 Which of the following products is formed when phenylamine (aniline) is reacted with **dilute** sulfuric acid?

- ☐ A 
- ☐ B 
- ☐ C 
- ☐ D 






(Total for Question 11 = 1 mark)

12 For the nitration of phenol, which is the most suitable set of conditions and the reason for its use?

|                          |   | Conditions                                                      | Reactivity of phenol to electrophiles compared with benzene |
|--------------------------|---|-----------------------------------------------------------------|-------------------------------------------------------------|
| <input type="checkbox"/> | A | dilute nitric acid at room temperature                          | more reactive                                               |
| <input type="checkbox"/> | B | concentrated nitric and sulfuric acid at room temperature       | more reactive                                               |
| <input type="checkbox"/> | C | concentrated nitric and sulfuric acid at 55 °C                  | the same                                                    |
| <input type="checkbox"/> | D | dilute nitric acid and dilute sulfuric acid at room temperature | less reactive                                               |

(Total for Question 12 = 1 mark)

13 Phenol reacts with excess bromine water to give as the organic product(s)





- ☐ A  and 
- ☐ B 
- ☐ C 
- ☐ D 

(Total for Question 13 = 1 mark)

14 An organic compound, **X**, shows the following properties:

- Oxidation of compound **X** produces a substance that reacts with 2,4-dinitrophenylhydrazine to give a yellow precipitate but does **not** react with Fehling's or Benedict's solution.
- Compound **X** reacts with ice-cold nitrous acid to form a compound that gives a yellow precipitate with an alkaline solution of phenol.

What is the formula of compound **X**?



(Total for Question 14 = 1 mark)

15 Which sequence shows the bases in order of decreasing strength?

- ☐ A  $\text{C}_6\text{H}_5\text{NH}_2 > \text{CH}_3\text{NH}_2 > \text{NH}_3$
- ☐ B  $\text{NH}_3 > \text{CH}_3\text{NH}_2 > \text{C}_6\text{H}_5\text{NH}_2$
- ☐ C  $\text{CH}_3\text{NH}_2 > \text{NH}_3 > \text{C}_6\text{H}_5\text{NH}_2$
- ☐ D  $\text{NH}_3 > \text{C}_6\text{H}_5\text{NH}_2 > \text{CH}_3\text{NH}_2$

(Total for Question 15 = 1 mark)

16 Bromoethane can be made by heating ethanol under reflux with 50% sulfuric acid and sodium bromide. When the mixture is distilled, the products include sulfur dioxide, bromine, hydrogen bromide and water as well as bromoethane.

The product mixture is shaken with sodium carbonate solution and later with anhydrous sodium sulfate before being re-distilled. Which of the following shows the correct list of impurities removed at each step?

|                          |   | Aqueous sodium carbonate wash       | Addition of sodium sulfate            |
|--------------------------|---|-------------------------------------|---------------------------------------|
| <input type="checkbox"/> | A | HBr                                 | $\text{SO}_2$ , $\text{Br}_2$ , water |
| <input type="checkbox"/> | B | $\text{SO}_2$ , $\text{Br}_2$       | HBr, water                            |
| <input type="checkbox"/> | C | $\text{SO}_2$ , HBr                 | $\text{Br}_2$ , water                 |
| <input type="checkbox"/> | D | $\text{SO}_2$ , $\text{Br}_2$ , HBr | water                                 |


(Total for Question 16 = 1 mark)

17 A compound is known to have either the structure  $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$  or  $\text{H}_2\text{NCH}_2\text{COOH}$ . Which of the following tests would best distinguish between the two compounds?

- ☐ A Reaction with concentrated aqueous sodium hydroxide.
- ☐ B Reaction with nitrous acid.
- ☐ C Reaction with aqueous sodium hydrogencarbonate.
- ☐ D Reaction with ethanoyl chloride.

(Total for Question 17 = 1 mark)

- 18 Poly(ethenol) is a water-soluble polymer. A section of the chain has the structure shown below.



The polymer is used for making hospital laundry bags so that laundry can be loaded directly into washing machines without it having to be handled.

Poly(ethenol) is water soluble because the polymer

- ☐ A is broken down by the water into monomers.
- ☐ B is broken down by the washing detergent.
- ☐ C breaks into monomers at the temperature of the wash.
- ☐ D forms many strong hydrogen bonds with the water.

(Total for Question 18 = 1 mark)

- 19 Which of the following substances is capable of damaging the ozone layer?

- ☐ A NaCl
- ☐ B CO<sub>2</sub>
- ☐ C C<sub>2</sub>HF<sub>5</sub>
- ☐ D C<sub>2</sub>F<sub>3</sub>Cl<sub>3</sub>

(Total for Question 19 = 1 mark)

- 20 Analysis suggests that a particular organic synthesis produces a medicine that contains trace impurities that may be hazardous. What is the best way for this discovery to be reported and evaluated?

- ☐ A In a scientific journal which subjects its articles to peer review.
- ☐ B On the Internet in an article on a website.
- ☐ C In a newspaper article in several broadsheet newspapers.
- ☐ D In a widely circulated magazine.

(Total for Question 20 = 1 mark)

**TOTAL FOR SECTION A = 20 MARKS**

## SECTION B

- 21 (a) Chromium is a typical transition metal, although its electronic configuration does **not** fit the general trend found in the first transition series.


Complete the electronic configurations in *s,p,d* notation for vanadium and chromium.

(1)

Vanadium: [Ar] .....

Chromium: [Ar] .....

- (b) Some interconversions found in the chemistry of chromium are shown below. Use this information to answer the questions that follow.



- (i) State **two** typical properties of transition metals, other than the formation of coloured ions, which are shown in the diagram above.

(2)

.....

.....

.....

- (ii) Use  $E^\ominus$  values from your data booklet to suggest a metal that could be used for **step 1**. Justify your answer by calculating  $E^\ominus$  for your cell.

(2)

- (iii) Explain, using oxidation numbers, whether or not the conversion in **step 3** is a redox reaction.

(2)


- (iv) The organic compound  $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$  that is used in **step 4** is 1,2-diaminoethane, often called ethylenediamine. It is a **bidentate ligand**. Explain the meaning of this term.

(1)

- (v) Explain, in terms of its structure, how  $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$  can act as a bidentate ligand whereas  $\text{H}_2\text{NNH}_2$  cannot.

(2)

(c) The half-equations relating the interconversion of the species  $\text{Cr}^{2+}(\text{aq})$ ,  $\text{Cr}^{3+}(\text{aq})$  and  $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$  are given below.



(i) Use your data booklet to find  $E^\ominus$  for each of the above half-equations.

(1)

**Half-equation I** ..... Volts

**Half-equation II** ..... Volts

\*(ii) Write the overall equation for the disproportionation of  $\text{Cr}^{3+}$  into  $\text{Cr}^{2+}$  and  $\text{Cr}_2\text{O}_7^{2-}$ .

Use the  $E^\ominus$  values you have obtained in (c)(i) to show whether or not this disproportionation is feasible under standard conditions.

(4)

.....

.....

.....

.....

.....

(Total for Question 21 = 15 marks)

22 (a) Tiglic acid is a compound that is used as a defensive agent by some beetles.

- (i) Tiglic acid contains, by mass, 60% carbon, 8% hydrogen, with the remainder being oxygen. Show that these data are consistent with the formula  $C_5H_8O_2$ .

(1)

- (ii) Tiglic acid contains a carbon-carbon double bond and a carboxylic acid group.

Suggest **one** test for each of these groups in tiglic acid. State what you would do and what you would see as a positive result for the tests.

(4)


Test for  $C=C$  .....

.....  
.....  
.....


Test for  $COOH$  .....

.....  
.....  
.....

(b) It is suggested that the structure of tiglic acid is either that of **A** or **B**.



**A**



**B**

- (i) State, with a reason, whether **B** is the *E*- or *Z*- isomer.

(2)

- (ii) The mass spectrum of tiglic acid shows two prominent peaks at mass/charge ratios 45 and 55. Write the formulae of the fragments giving rise to each of these peaks.

(2)


45 .....

55 .....

- (iii) Does this data from the mass spectrum **alone** enable you to decide which of **A** or **B** is the structure of tiglic acid? Explain your answer.

(1)

- (c) The position of a C=C double bond in a molecule can be determined by ozonolysis. The compound is reacted with ozone and then dilute acid, two carbonyl compounds being produced as shown below.



Ozonolysis of tiglic acid gives two carbonyl compounds, **C** and **D**.

Compound **C** gives a silver mirror with Tollens' reagent and gives iodoform with iodine in alkali.

Compound **D** does **not** give a silver mirror with Tollens' reagent, but does give iodoform with iodine in alkali.

- \*(i) From the results of the experiments, deduce the functional groups present in **C** and **D**. By considering the two possible structures for tiglic acid, give the structural formulae of **C** and **D**.


From the structures you have drawn, state which of the structures **A** or **B** could represent tiglic acid.

(6)


- (ii) Explain whether or not these tests show definitely that your answer to (c)(i) represents tiglic acid.

(1)

(d) Lactic acid is a chiral molecule that is found in sweat as the (–) isomer only. Its structural formula is



(i) Lactic acid can be made from ethanol in three steps.



Give the structural formula of the intermediate **X** and the reagents and conditions required for **steps 1** and **2**.

(4)

Step 1 .....

.....

Step 2 .....

.....

(ii) Classify the type and mechanism of the reaction that occurs in **step 2**.

(1)

.....

\*(iii) By considering the stereochemistry of the mechanism in **step 2**, explain why this synthesis would **not** give a single optical isomer of lactic acid.

(2)

.....

.....

.....

(iv) Suggest why synthetic pathways for the manufacture of pharmaceuticals may require reactions that are highly stereospecific.

(1)


.....

.....

.....

(Total for Question 22 = 25 marks)

23 Proteins are polymers of  $\alpha$ -amino acids, the simplest of which is glycine.



(a) (i) Draw the **structural** formula for the zwitterion of glycine in the solid state. (1)

(ii) Explain, on the basis of your answer to (a)(i), why glycine has a relatively high melting temperature for such a small molecule. (2)

.....

.....

.....

.....

(iii) Draw the structure of the protein chain that would be formed if glycine alone were to be polymerized. Show part of the chain containing two glycine residues. (2)

(5)

---

## SECTION C


24 Read the passage below carefully and answer the questions which follow.

**Molecular structure and colour chemistry**


The sight of tubes of paint or of coloured pencils in an artists' supply shop is something that most people enjoy; we love colour.

The ability to synthesise brightly-coloured compounds coincides with the rapid growth of the organic chemicals industry. Synthetic organic dyes started to appear in the mid-19th century when William Perkin synthesised Mauve in 1856 at the age of 18. He was trying to synthesise quinine even though he did not know the structure of the molecule.

In the 19th century many chemists did not believe that molecules existed. The work of Butlerov, Couper, and notably Kekulé showed that molecules not only exist but have specific structures. In 1865 Kekulé suggested a ring structure for the aromatic compound benzene which he represented as



Kekulé knew that benzene does not react with bromine water. Later work showed that the enthalpy change of hydrogenation of the compound is  $-205 \text{ kJ mol}^{-1}$ , rather than the value of  $-360 \text{ kJ mol}^{-1}$  that would be expected if the structure was exactly as shown above, given that the enthalpy change of hydrogenation for cyclohexene to cyclohexane



is  $-120 \text{ kJ mol}^{-1}$ .

When Greiss in 1856 discovered diazotisation and the azo dyes, he used a reaction characteristic of aromatic amines. Witt, in 1876, found the functional groups in the dye molecule that make it water-soluble and enable it to attach to the cloth fibres. Graebe, Liebermann and Perkin in 1869 patented the synthesis of alizarin, found in madder root grown in Holland and Von Baeyer synthesised indigo in 1880, until then grown in India. Synthetic dyes were made available in large quantities and were cheaper than the sources from plants.

Now the organic chemical industry produces a vast range of pigments and dyestuffs for use in paints and for fabrics, inks and other materials, making our world the most colourful that it has ever been.

(a) (i) Explain why Perkin's attempted synthesis of quinine was almost certain to fail.

(1)

(ii) Suggest the effect that the growth of the organic chemicals industry in the late 19th century had on Holland and India in particular.

(2)


(b) (i) What observation did Kekulé make to show that benzene does **not** react with bromine water? Explain the significance of this with reference to his representation of the molecule.

(2)

(ii) Explain, in terms of the bonding in the benzene ring, why the enthalpy of hydrogenation is less exothermic than would be expected from a molecule with three double bonds.

(3)

(c) The first steps in the preparation of an azo dye from benzene are shown below.



Give the mechanism for the reaction in **step 1**, including the equation for the formation of the electrophile.

(4)

Equation for formation of electrophile

Mechanism

- (d) (i) Phenylamine is converted into the benzenediazonium cation using sodium nitrite and hydrochloric acid at a temperature between 0 °C and 10 °C.

Explain why the temperature must **not** be lower or higher than these limits if a good yield is to be obtained.

(2)

.....

.....

.....

.....


- (ii) Draw the structural formula of the benzenediazonium cation showing all the bonds and the charge.

(1)

- (iii) Suggest how you could convert a sample of the benzenediazonium cation into an azo dye. Give the name of the other compound you would use and the skeletal formula of the azo dye you would obtain.

(3)

(e) The structural formula of methyl orange is given below.



Suggest the main features of methyl orange which make it water-soluble, giving your reasons.

(2)

---

(Total for Question 24 = 20 marks)

---

---

**TOTAL FOR SECTION C = 20 MARKS**  
**TOTAL FOR PAPER = 90 MARKS**

mock papers 5

SECTION A

1 The acid used in a standard hydrogen electrode to provide a  $1 \text{ mol dm}^{-3}$  solution of hydrogen ions is

- ☐ A ethanoic acid.
- ☐ B phosphoric(V) acid.
- ☐ C sulfuric acid.
- ☐ D hydrochloric acid.

---

(Total for Question 1 = 1 mark)

2 To measure the standard electrode potential for the  $\text{Ag}^+(\text{aq})|\text{Ag}(\text{s})$  electrode, the most suitable chemical for the solution in a salt bridge to connect the two half cells is

- ☐ A potassium chloride.
- ☐ B potassium iodide.
- ☐ C potassium nitrate.
- ☐ D potassium sulfate.

---

(Total for Question 2 = 1 mark)

3 For any reversible reaction,  $\ln K$  is proportional to

- ☐ A both  $\Delta S_{\text{total}}$  and  $E_{\text{cell}}$ .
- ☐ B  $\Delta S_{\text{total}}$  but not  $E_{\text{cell}}$ .
- ☐ C  $E_{\text{cell}}$  but not  $\Delta S_{\text{total}}$ .
- ☐ D neither  $E_{\text{cell}}$  nor  $\Delta S_{\text{total}}$ .

---

(Total for Question 3 = 1 mark)

- 4 This question is about a titration to determine the iron content of a tablet. The iron(II) ions in the tablet are oxidized to iron(III) ions by acidified manganate(VII) ions which are reduced to manganese(II) ions.

(a) The mole ratio of iron(II) to manganate(VII) ions in the reaction is

(1)

|                            | $\text{Fe}^{2+}$ | $\text{MnO}_4^-$ |
|----------------------------|------------------|------------------|
| <input type="checkbox"/> A | 1                | 5                |
| <input type="checkbox"/> B | 2                | 5                |
| <input type="checkbox"/> C | 5                | 2                |
| <input type="checkbox"/> D | 5                | 1                |

- (b) A 0.200 g tablet is dissolved to make exactly 100 cm<sup>3</sup> of solution. 10 cm<sup>3</sup> of this solution is found to contain  $5.38 \times 10^{-5}$  mol of iron(II) ions.

The percentage by mass of iron ( $A_r = 55.8$ ) in the tablet is

(1)

☐ A  $\frac{5.38 \times 10^{-5} \times 55.8}{0.200} \times \frac{100}{10} \times 100\%$


☐ B  $\frac{5.38 \times 10^{-5}}{55.8 \times 0.200} \times \frac{100}{10} \times 100\%$

☐ C  $\frac{5.38 \times 10^{-5} \times 55.8}{0.200} \times \frac{10}{100} \times 100\%$

☐ D  $\frac{5.38 \times 10^{-5} \times 0.200}{55.8} \times \frac{10}{100} \times 100\%$





(Total for Question 4 = 2 marks)

- 5 Some fuel cells are used to produce energy from the combination of hydrogen and oxygen. The half-equations for a hydrogen-oxygen fuel cell in alkaline solution are





(a) The equation for the cell reaction is

(1)



(b) From the data above,  $E_{\text{cell}}$  for the reaction in the hydrogen-oxygen fuel cell is

(1)



(c) An advantage of operating a hydrogen fuel cell over a conventional jet engine for powering an aeroplane is

(1)

☐ A it is 100% efficient.


☐ B it does not produce greenhouse gas emissions.

☐ C it does not produce emissions which damage the ozone layer.

☐ D hydrogen is easier to transport.

(Total for Question 5 = 3 marks)

- 6 Benzocaine is used as a local anaesthetic.



Separate samples of a solution of benzocaine are added to 2,4-dinitrophenylhydrazine, hot aqueous sodium hydroxide, and dilute hydrochloric acid.

Which chemicals react with benzocaine?

- ☐ A All three
- ☐ B Only sodium hydroxide and hydrochloric acid
- ☐ C Only hydrochloric acid
- ☐ D Only sodium hydroxide

(Total for Question 6 = 1 mark)

- 7 In the first commonly used breathalyser, acidified potassium dichromate(VI) was used which was reduced to chromium(III) by alcohol.

The colour change seen when alcohol was present in a motorist's breath is from

- ☐ A orange to green.
- ☐ B orange to yellow.
- ☐ C yellow to green.
- ☐ D yellow to orange.


(Total for Question 7 = 1 mark)

- 8 For which of the following are both tests acceptable evidence for a driver to be successfully prosecuted for excess alcohol in the blood in many countries?

- ☐ A A dichromate(VI) breath analyser and a blood test.
- ☐ B A dichromate(VI) breath analyser and a fuel cell breath analyser.
- ☐ C A dichromate(VI) breath analyser and an infrared breath analyser.
- ☐ D An infrared breath analyser and a blood test.

(Total for Question 8 = 1 mark)

9 The structural formula of caffeine,  $C_8H_{10}O_2N_4$ , is shown below.



(a) How many main peaks would you expect in the proton nuclear magnetic resonance spectrum of caffeine?

(1)

- ☐ A 1
- ☐ B 2
- ☐ C 3
- ☐ D 4

(b) At which of the following wavenumbers is an absorption peak **not** present in the infrared spectrum of caffeine?

(1)

- ☐ A  $3600\text{ cm}^{-1}$
- ☐ B  $2925\text{ cm}^{-1}$
- ☐ C  $1690\text{ cm}^{-1}$
- ☐ D  $1660\text{ cm}^{-1}$

(c) The parent ion peak of caffeine in the mass spectrum of caffeine would be at m/e ratio

(1)

- ☐ A 101
- ☐ B 102
- ☐ C 193
- ☐ D 194

(d) Intermolecular forces between caffeine molecules would include

(1)

- ☐ A London forces only.
- ☐ B London forces and hydrogen bonds.
- ☐ C London forces and permanent dipole forces.
- ☐ D London forces, permanent dipole forces, and hydrogen bonds.

(Total for Question 9 = 4 marks)

10 (a) The principal advantage of combinatorial chemistry over traditional methods for developing pharmaceuticals is that

(1)

- ☐ A very small amounts of compounds are used.
- ☐ B many more compounds can be made in a given time.
- ☐ C compounds formed are more powerful drugs.
- ☐ D compounds formed have fewer side effects.




(b) A refinement of combinatorial chemistry involves initially attaching compounds firmly to polymer beads. The type of attachment is most likely to be by

(1)

- ☐ A metallic bonds.
- ☐ B covalent bonds.
- ☐ C London forces.
- ☐ D permanent dipole forces.

(Total for Question 10 = 2 marks)

11 Four reactions involving the transition elements copper and chromium are given below.



(a) Which reaction produces a dark blue solution?

(1)

☐ A 1

☐ B 2

☐ C 3

☐ D 4

(b) Which two reactions show the amphoteric behaviour of a metal hydroxide?

(1)

☐ A 1 and 2

☐ B 2 and 3

☐ C 2 and 4

☐ D 3 and 4

(c) Predict, without calculation, which reaction has the most negative value for  $\Delta S_{\text{system}}$ .

(1)

☐ A 1

☐ B 2

☐ C 3

☐ D 4


(Total for Question 11 = 3 marks)

**TOTAL FOR SECTION A = 20 MARKS**

## SECTION B

12 This question is about benzene and its compounds.

- (a) A Kekulé structure of benzene suggests the molecule consists of alternate double and single carbon to carbon bonds.



The standard enthalpy change of hydrogenation of a carbon to carbon double bond is  $-120 \text{ kJ mol}^{-1}$ .

- (i) Calculate the standard enthalpy change of hydrogenation of benzene for the Kekulé structure.

(1)

- \*(ii) The actual standard enthalpy change of hydrogenation of benzene is  $-208 \text{ kJ mol}^{-1}$ .


Use this information and your answer to (i), to calculate the difference in stability of benzene and the Kekulé structure.

What does this tell us about the bonding in benzene?

Explain how this influences the type of chemical reactions that benzene undergoes.

(4)

(b) The flow chart shows some typical reactions of benzene.



(i) Identify the two chemicals needed to make nitrobenzene from benzene.

(2)

(ii) Classify the type of reagent required in all three reactions.

(1)

(iii) Show, with appropriate diagrams, the mechanism for the reaction of benzene with bromine, in the presence of iron(III) bromide, to form bromobenzene. Include the formation of the ion attacking the benzene molecule.

(4)

(iv) Give the structural formula and the name for compound A.

(2)

Name.....

(c) (i) Explain why phenol,  $\text{C}_6\text{H}_5\text{OH}$ , and methoxybenzene,  $\text{C}_6\text{H}_5\text{OCH}_3$ , are much more reactive than benzene with bromine.

(3)

.....

.....

.....

.....

.....

.....

(ii) Write the equation for the reaction between phenol and bromine water. State symbols are **not** required.

(2)

(d) Diphenylmethane,  $(\text{C}_6\text{H}_5)_2\text{CH}_2$ , is often added to soap to create the pleasant smell of geraniums. It can be made from benzene and one other reagent, using a suitable catalyst.

Identify the other starting material and the catalyst by name or formula.

(2)

.....

.....


(Total for Question 12 = 21 marks)

13 This question is about 1,2-diaminoethane,  $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$ .

- (a) Write an equation for the reaction between 1,2-diaminoethane and **excess** dilute hydrochloric acid.

(2)

- (b) When an aqueous solution of nickel(II) ions is added to 1,2-diaminoethane, a complex ion forms.



- (i) Suggest the colour of this complex.

(1)

- (ii) Without using the data booklet, suggest why the complex formed is more stable than  $\text{Ni}(\text{H}_2\text{O})_6^{2+}$  by considering the appropriate entropy change.

(2)

- (iii) This complex can exist as two structures, which are non-superimposable mirror images.

Suggest the physical property that would enable you to distinguish between these two structures.

(1)

(c) 1,2-diaminoethane reacts with butane-1,2-dioyl dichloride,  $\text{ClOCCH}_2\text{CH}_2\text{COCl}$ , to form a polymer.

(i) Draw a displayed formula for this polymer showing one repeat unit.

(2)

(ii) Give, and explain, the name for this type of polymerization reaction.

(2)

\*(iii) State and explain the intermolecular forces between two of these polymer chains.

(5)

(Total for Question 13 = 15 marks)

14 An organic compound **X** contains carbon, hydrogen, oxygen and nitrogen.

0.132 g of **X** is burned completely in oxygen to produce 0.072 g of water, 0.176 g of carbon dioxide and 24.0 cm<sup>3</sup> of nitrogen.

[Molar volume of nitrogen under the conditions of the experiment = 24 000 cm<sup>3</sup> mol<sup>-1</sup>]

(a) Calculate the empirical formula of compound **X**.

(5)

(b) The molar mass of **X** is 132 g mol<sup>-1</sup>. Deduce the molecular formula for **X**. Show how you arrived at your answer.

(1)

- (c) When **X** is refluxed with concentrated hydrochloric acid for several hours, cooled and neutralized, there is only one organic product, **Y**, which has the molecular formula  $C_2H_5O_2N$ .

One mole of **Y** will react with either one mole of hydrochloric acid or one mole of sodium hydroxide solution.

When **Y** is sprayed with a solution of ninhydrin and heated, a purple colour is observed.

- (i) Use **all** the information above to deduce the functional groups present in **Y** and to classify the type of compound it is. Justify your answer.

(3)

.....

.....

.....

.....

.....

.....

- (ii) Deduce the displayed formula for **Y** and give its name.

(2)

Name of **Y**.....

- (iii) Hence give the structural formula for **X**.

(1)

(Total for Question 14 = 12 marks)

**TOTAL FOR SECTION B = 48 MARKS**

## SECTION C

15


**Some unusual oxidation states**

You will probably have seen manganese in its common oxidation states during your study of chemistry. Manganese(II) sulfate occurs as very pale pink crystals in the hydrated form. Manganese(IV) oxide is a black powder which is often used as a catalyst. Potassium manganate(VII) occurs as very dark purple crystals and forms a purple aqueous solution, which is a powerful oxidizing agent.

You are less likely to have seen compounds containing the other oxidation states of manganese, which are manganese(VI), manganese(V), manganese(III) and manganese(I). However, compounds containing each of these four oxidation states can be prepared.

**Manganese(VI)**


Manganese(VI), in  $\text{MnO}_4^{2-}$ , can be prepared in a reverse disproportionation reaction, by reacting manganate(VII) ions with manganese(IV) oxide in alkali.

**Equation 1**

The reaction is not thermodynamically favourable under standard conditions. However, the  $E_{\text{cell}}$  value can be made positive by increasing the concentration of hydroxide ions so that green manganate(VI) ions form.

**Manganese(V)**

Manganese(V) can be formed by adding a little potassium manganate(VII) to very concentrated ( $12 \text{ mol dm}^{-3}$ ) aqueous sodium hydroxide. The solution slowly becomes blue as manganate(V) ions,  $\text{MnO}_3^-(\text{aq})$ , form. The ionic half-equations are:

**Equation 2****Equation 3****Manganese(III)**

A deep red solution containing manganese(III) ions is formed by the oxidation of manganese(II) hydroxide by potassium manganate(VII) in acid solution. The ionic equation for the reaction is:

**Equation 4****Manganese(I)**

Manganese(I) ions are not stable in aqueous solution, but do form stable complex ions. They can be made by reducing hexacyanomanganate(II) ions,  $\text{Mn}(\text{CN})_6^{4-}$ , to hexacyanomanganate(I) ions,  $\text{Mn}(\text{CN})_6^{5-}$ .

(a) (i) Give the formula of manganese(IV) oxide.

(1)

(ii) How do catalysts speed up reactions?

(2)


(iii) Explain how transition metal ions can act as homogeneous catalysts.

(2)

(b) (i) Suggest why the preparation of manganate(VI) ions,  $\text{MnO}_4^{2-}$ , in **equation 1**, may be described as a reverse disproportionation reaction by considering the relevant oxidation states.

(2)

- (ii) The two half-equations which are combined to form **equation 1** are



Explain, by reference to these half-equations, why increasing the concentration of hydroxide ions alters the electrode potential to make the preparation of manganate(VI) ions more likely.

(3)

.....

.....

.....

.....

- (c) Use equations **2** and **3** to answer the following questions.

- (i) Identify the gas formed in the preparation of manganate(V) ions.

(1)

.....

- (ii) By appropriately combining these two equations, write the ionic equation for the formation of manganate(V) ions from manganate(VII) ions.

(2)

- (iii) Identify the main hazard and state how you would minimize the associated risk in this preparation of manganate(V) ions.

(2)

.....

.....

.....

.....

(d) Identify the reagents you would use to make manganese(II) hydroxide for the preparation of manganese(III) ions.

(2)

(e) (i) Draw a dot and cross diagram to show the electron arrangement in the cyanide ion,  $\text{CN}^-$ .

(2)

(ii) Explain how the cyanide ion acts as a ligand.

(2)

(iii) Suggest the name of the shape of the hexacyanomanganate(I) ion.

(1)

---

(Total for Question 15 = 22 marks)

---

**TOTAL FOR SECTION C = 22 MARKS**

**TOTAL FOR PAPER = 90 MARKS**